



**Ontario Health**  
Cancer Care Ontario

Guideline MOTAC-4 Version 2 IN REVIEW

A Quality Initiative of the  
Program in Evidence-Based Care (PEBC), Ontario Health (Cancer Care  
Ontario)

## Clinical Utility of Multigene Profiling Assays in Early-Stage Invasive Breast Cancer

*P. Blanchette, D. Sivajohanathan, J. Bartlett, A. Eisen, H. Feilotter, R. Pezo, G. Turashvili,  
P. Williams, and the Multigene Profiling Assays in Early-Stage Invasive Breast Cancer Expert  
Panel*

Report Date: January 28, 2022

An assessment conducted in February 2025 placed Guideline MOTAC-4 Version 2 IN REVIEW. This means that it is undergoing a review for currency and relevance. It is still appropriate for this document to be available while this updating process unfolds. The PEBC has a formal and standardized process to ensure the currency of each document ([PEBC Assessment & Review Protocol](#))

Guideline MOTAC-4 Version 2 is comprised of 5 sections. You can access the summary and full report here:

<https://www.cancercareontario.ca/en/guidelines-advice/types-of-cancer/31766>

|            |                              |
|------------|------------------------------|
| Section 1: | Recommendations Summary      |
| Section 2: | Guideline                    |
| Section 3: | Guideline Methods Overview   |
| Section 4: | Evidence Review              |
| Section 5: | Internal and External Review |

For information about this document, please contact Dr. Phillip Blanchette, the lead author, through the PEBC at:  
Phone: 905-527-4322 ext. 42822 Fax: 905-526-6775 E-mail: [ccopgi@mcmaster.ca](mailto:ccopgi@mcmaster.ca)

For information about the PEBC and the most current version of all reports, please visit the OH (CCO) website at <https://www.cancercareontario.ca/en/guidelines-advice> or contact the

PEBC office at:

Phone: 905-527-4322 ext. 42822 Fax: 905-526-6775 E-mail: [ccopgi@mcmaster.ca](mailto:ccopgi@mcmaster.ca)

**PEBC Report Citation (Vancouver Style):** Blanchette P, Sivajohanathan D, Bartlett J, Eisen A, Feilotter H, Pezo R, et al. Clinical utility of multigene profiling assays in early-stage invasive breast cancer. Toronto (ON): Ontario Health (Cancer Care Ontario); 2022 *January 28* [In Review 2025 Feb]. Program in Evidence-Based Care Guideline No.: MOTAC-4 Version 2 IN REVIEW.

#### **PUBLICATIONS RELATED TO THIS REPORT**

Blanchette P, Sivajohanathan D, Bartlett J, Eisen A, Feilotter H, Pezo R, et al. Clinical Utility of Multigene Profiling Assays in Early-Stage Invasive Breast Cancer: An Ontario Health (Cancer Care Ontario) Clinical Practice Guideline. *Curr Oncol.* 2022;29:2599-2615.

#### *Copyright*

This report is copyrighted by Ontario Health (Cancer Care Ontario); the report and the illustrations herein may not be reproduced without the express written permission of Ontario Health (Cancer Care Ontario). Ontario Health (Cancer Care Ontario) reserves the right at any time, and at its sole discretion, to change or revoke this authorization.

#### *Disclaimer*

Care has been taken in the preparation of the information contained in this report. Nevertheless, any person seeking to consult the report or apply its recommendations is expected to use independent medical judgment in the context of individual clinical circumstances or to seek out the supervision of a qualified clinician. Ontario Health (Cancer Care Ontario) makes no representations or guarantees of any kind whatsoever regarding the report content or its use or application and disclaims any responsibility for its use or application in any way.

# Clinical Utility of Multigene Profiling Assays in Early-Stage Invasive Breast Cancer

## Recommendations

*This is a quick reference guide and provides the guideline recommendations only. For key evidence associated with each recommendation, the systematic review, and the guideline development process, see the Full Report.*

### GUIDELINE OBJECTIVES

To update clinical guidance on the use of multigene profiling assays in individuals with early-stage invasive breast cancer.

### TARGET POPULATION

Individuals diagnosed with early-stage invasive breast cancer for whom further information is needed for prognosis and treatment decision making. In this guideline, early-stage invasive breast cancer is defined as stage I to III breast cancers that are surgically operable and do not have evidence of inflammatory, locally recurrent or distant metastatic disease with pT1-T3, pN0-N1a based on surgical pathologic staging.

### INTENDED USERS

This guideline is targeted for clinicians and policy makers involved in the diagnosis and treatment of breast cancer.

### PREAMBLE

The purpose of this guideline is to determine the clinical utility of multigene profiling assays (i.e., Oncotype DX, MammaPrint, Prosigna, EndoPredict, and Breast Cancer Index), not to identify which assay is better. No prospective studies have compared these head-to-head. Given that the assays use different scoring systems and classification systems, please refer to Table 1-1 for a summary of each of the assays. Further, this guideline does not cover the utility of multigene profiling assays in helping to guide clinical treatment decisions regarding the use of either neoadjuvant chemotherapy or radiation.

### RECOMMENDATIONS, KEY EVIDENCE, AND JUSTIFICATION

|   |
|---|
| <b>Recommendation 1</b>   |
| In patients with early-stage estrogen receptor (ER)-positive/human epidermal growth factor 2 (HER2)-negative breast cancer, clinicians should consider using multigene profiling assays (i.e., Oncotype DX, MammaPrint, Prosigna, EndoPredict, and Breast Cancer Index) to help guide the use of systemic therapy.  |
| <b>Qualifying Statements for Recommendation 1</b>   |
| <ul style="list-style-type: none"> <li>• There is currently insufficient evidence to use multigene profiling assays among patients with either HER2-positive or triple negative breast cancers.</li> <li>• Multigene profiling assays are recommended for use in patients with lymph node-negative or lymph node-positive (1-3 lymph nodes) disease who are under consideration for adjuvant chemotherapy if the use is supported by other clinical, pathological, or patient-related factors. Clinical and pathological features include patient age, tumour grade, tumour size and nodal status.</li> </ul> |

- One multigene profiling assay should be requested per patient to guide a specific treatment decision. Requesting multiple tests with different multigene profiling assays on an individual tumour specimen to guide a single treatment decision is discouraged. Additional testing may be considered for patients with either repeat metachronous breast cancer diagnoses or synchronous breast cancer diagnoses where tumour specimens display varying morphologies, grade or hormone receptor status.
- Multigene profiling assays should be interpreted cautiously in premenopausal patients where a significant benefit from adjuvant chemotherapy may still exist despite a low-risk score.

**Recommendation 2**

In patients with early-stage node-negative ER-positive/HER2-negative disease, clinicians may use a low-risk result from Oncotype DX, MammaPrint, Prosigna, EndoPredict/EPclin, or Breast Cancer Index assays to support a decision not to use adjuvant chemotherapy.

**Qualifying Statements for Recommendation 2**

- Patients <50 years of age may still benefit from chemotherapy despite low-risk scores from multigene assay testing. Risk scores should be interpreted with caution and decisions should be made while considering other clinical, pathological, or patient-related factors.
- Treatment decisions should be based on all available clinical and pathological information for each patient, rather than depending entirely on multigene profiling test results.
- In patients with a low-grade tumour (i.e., grade 1) less than 1 cm in size, the Working Group members do not recommend a multigene assay profiling as this is unlikely to inform a treatment decision to use adjuvant chemotherapy.

**Recommendation 3**

In patients with node-negative ER-positive/HER2-negative disease, clinicians may use a high-risk result from Oncotype DX to support a decision to offer chemotherapy. A high Oncotype DX recurrence score is capable of predicting adjuvant chemotherapy benefit.

**Qualifying Statements for Recommendation 3**

- MammaPrint, Prosigna, EndoPredict or EPclin, and Breast Cancer Index do not have sufficient evidence to support a predictive benefit of adjuvant chemotherapy among clinically low-risk patients with breast cancer whose multigene profiling testing indicates a high-risk score.

**Recommendation 4**

In postmenopausal patients with ER-positive/HER2-negative tumours and one to three nodes involved (N1a disease), clinicians may withhold chemotherapy based on a low-risk Oncotype DX or MammaPrint score if the decision is supported by other clinical, pathological, or patient-related factors.

**Qualifying Statements for Recommendation 4**

- Premenopausal patients <50 years of age have a significant benefit from chemotherapy despite low-risk scores from multigene assay testing. Risk scores should be interpreted with caution and decisions should be made while considering other clinical, pathological, or patient-related factors.
- It is uncertain whether at least some of the benefit of chemotherapy among premenopausal patients may be due to chemotherapy induced amenorrhea versus the cytotoxic effects of treatment.

- The Prosigna, EndoPredict/EPclin, and Breast Cancer Index assays can identify low-risk node-positive patients whose prognostic outcomes are favourable; however, these assays have not demonstrated predictive evidence to support withholding adjuvant chemotherapy among higher risk, node-positive, ER-positive, HER2-negative breast cancer patients.

**Recommendation 5**

The evidence to support the use of molecular profiling to select the duration of endocrine therapy is evolving. In patients with ER-positive disease, clinicians may consider using a Breast Cancer Index (BCI) (H/I) high assay result to support a decision to extend adjuvant endocrine therapy if the decision is supported by other clinical, pathological, or patient-related factors.

**Qualifying Statements for Recommendation 5**

- While a number of studies have demonstrated clinical utility of BCI for extending adjuvant endocrine therapy, the preliminary results of the NSABP B42 trial are negative leading to some uncertainty. Treatment decisions should be based on all available clinical and pathological information for each patient, rather than depending only on multigene profiling tests.
- MammaPrint, Oncotype DX, Prosigna, and EndoPredict currently have insufficient evidence to guide extension of adjuvant endocrine therapy; however, these molecular assays may prognosticate a very low rate of disease recurrence that might not justify an extension of endocrine therapy.

Figure 1-1. Multigene Profiling Assay Decision Tree for Adjuvant Chemotherapy in Node-Negative Patients




Figure 1-2. Multigene Profiling Assay Decision Tree for Adjuvant Chemotherapy in Node-Positive Patients




Figure 1-3. Multigene Profiling Assay Decision Tree for Extended Adjuvant Endocrine Therapy






Table 1-1. Summary of assay characteristics.

| Assay                              | Oncotype DX   | MammaPrint   | Prosigna  | EndoPredict   | Breast Cancer Index   |
|------------------------------------|---|--|---|---|---|
| Tissue Required                    | FFPE  | FFPE or fresh tissue   | FFPE  | FFPE  | FFPE  |
| Technique                          | qRT-PCR   | Microarray   | qRT-PCR and nCounter DX Analysis System   | qRT-PCR   | qRT-PCR   |
| Assay Output                       | RS (0-100)  | MammaPrint Index<br>Risk of distant recurrence at 5 years      | Intrinsic subtype and ROR score (0-100)   | EPclin score (1-6)<br>Molecular score (1-15)  | BCI score (0-10) and BCI (H/I) low and BCI (H/I) high (ratio HoxB13 and interleukin-17B receptor)   |
| Categories for Risk Measurement    | <i>TAILORx categories</i><br>Low: ≤15<br>Intermediate: 16-25<br>High: 26-100<br><br><i>Pre-TAILORx categories</i><br>Low: <18<br>Intermediate: 18-30<br>High: ≥31 | Low: 0 to 1<br>High: -1 to 0                                   | <i>LN-negative</i><br>Low: 0-40<br>Intermediate: 41-60<br>High: 61-100<br><br><i>LN-positive (1-3 nodes)</i><br>Low: 0-40<br>High: 41-100 | <i>EPclin score</i><br>Low: <3.3<br>High: ≥3.3<br><br><i>Molecular score</i><br>Low: <5<br>High: ≥5 | <i>BCI predictive H/I</i><br>Low: <0.06<br>High: ≥0.06<br><br><i>BCI prognostic node-negative</i><br>Low: <5.0825<br>Intermediate: 5.0825-6.5025<br>High: ≥6.5025<br><br><i>BCI prognostic node-positive</i><br>Low: <6.93<br>High: ≥6.93 |
| Regulatory Approval or Endorsement | Assay conducted in centralized Exact Science's CLIA-certified lab   | FDA cleared for Agendia centralized lab testing in FFPE (2015) | FDA cleared for decentralized testing (2014)  | CE Mark for decentralized testing (2012)  | Assay conducted in centralized CAP/CLIA-certified lab   |
| Manufacturer                       | Exact Sciences Corp.  | Agendia  | Veracyte  | Myriad Genetics, Inc.   | Biotheranostics, Inc.   |
| Testing Location                   | Central (1 laboratory in US)  | Central (1 laboratory in the Netherlands, 1 in US)             | Various labs across US, UK  | Central laboratory in the US  | Central (1 laboratory in US)  |
| Genes, n                           | 21-gene assay   | 70-gene assay  | 50-gene assay   | 12-gene assay<br>EPclin score: 12-gene assay plus tumour size and nodal status                      | <i>HOXB13:IL17BR</i> expression ratio (H/I) and Molecular Grade Index   |

**Abbreviations:** BCI (H/I), Breast Cancer Index (HOXB13/IL17BR); CAP: College of American Pathologists; CLIA: Clinical Laboratory Improvement Amendments; EPclin, EndoPredict clinical score; ER, estrogen receptor; FDA, Food and Drug Administration; FFPE, formalin-fixed paraffin-embedded; LN, lymph node; qRT-PCR, quantitative reverse-transcription polymerase chain reaction; ROR: risk of recurrence; RS, recurrence score; UK: United Kingdom; US, United States