



Guideline 17-12

A Quality Initiative of the
Program in Evidence-Based Care (PEBC), Cancer Care Ontario (CCO)

Indications for Hyperthermic Intraperitoneal Chemotherapy with Cytoreductive Surgery

R. Auer, D. Sivajohanathan, J. Biagi, J. Conner, E. Kennedy, T. May and the HIPEC Guideline Group

Report Date: October 30, 2019

An assessment conducted in January 2026 deferred the review of Guideline 17-12. This means that the document remains current until it is assessed again next year. The PEBC has a formal and standardized process to ensure the currency of each document ([PEBC Assessment & Review Protocol](#))

Guideline 17-12 is comprised of 5 sections. You can access the summary and full report here:

<https://www.cancercareontario.ca/en/guidelines-advice/types-of-cancer/61856>

Section 1:	Recommendations
Section 2:	Recommendations and Key Evidence
Section 3:	Guideline Methods Overview
Section 4:	Systematic Review
Section 5:	Internal and External Review

For information about this document, please contact Dr. Rebecca Auer, the corresponding author, through the PEBC via:
Phone: 905-527-4322 ext. 42822 Fax: 905 526-6775 E-mail: ccopgi@mcmaster.ca

For information about the PEBC and the most current version of all reports, please visit the CCO website at <http://www.cancercareontario.ca/en/guidelines-advice> or contact the PEBC office at:

Phone: 905-527-4322 ext. 42822 Fax: 905 526-6775 E-mail: ccopgi@mcmaster.ca

PEBC Report Citation (Vancouver Style): Auer R, Sivajohanathan D, Biagi J, Connor J, Kennedy E, May T et al. Indications for cytoreductive surgery with hyperthermic intraperitoneal chemotherapy. Toronto (ON): Cancer Care Ontario; 2019 October 30. Program in Evidence-Based Care Guideline No.: 17-12.

Copyright

This report is copyrighted by Cancer Care Ontario; the report and the illustrations herein may not be reproduced without the express written permission of Cancer Care Ontario. Cancer Care Ontario reserves the right at any time, and at its sole discretion, to change or revoke this authorization.

Disclaimer

Care has been taken in the preparation of the information contained in this report. Nevertheless, any person seeking to consult the report or apply its recommendations is expected to use independent medical judgment in the context of individual clinical circumstances or to seek out the supervision of a qualified clinician. Cancer Care Ontario makes no representations or guarantees of any kind whatsoever regarding the report content or its use or application and disclaims any responsibility for its use or application in any way.

Indications for Hyperthermic Intraperitoneal Chemotherapy with Cytoreductive Surgery

Recommendations

This is a quick reference guide and provides the guideline recommendations only. For key evidence associated with each recommendation, the systematic review, and the guideline development process, see the Full Report.

GUIDELINE OBJECTIVES

To determine evidence-based indications for hyperthermic intraperitoneal chemotherapy (HIPEC) with cytoreductive surgery (CRS).

TARGET POPULATION

Adults (≥ 18 years old) with a diagnosis of mesothelioma, appendiceal (including appendiceal mucinous neoplasm), colorectal, gastric, ovarian, or primary peritoneal carcinoma.

INTENDED USERS

This guideline is intended for clinicians involved in the care of patients with mesothelioma, appendiceal (including appendiceal mucinous neoplasm), colorectal, gastric, ovarian, or primary peritoneal carcinoma.

RECOMMENDATIONS, KEY EVIDENCE, AND INTERPRETATION OF EVIDENCE

NOTE: This guideline addresses the role of HIPEC with CRS and not the role of CRS alone. Interventions and terms are reported as stated in the individual papers. While there is a lack of evidence to make recommendations for many of the target sites, it is noted that there are a large number of ongoing randomized controlled trials (RCTs). This guideline will be reviewed annually for any new evidence. When writing these recommendations, the Working Group considered overall survival (OS) to be a critical outcome, and progression-free survival (PFS), recurrence-free survival (RFS), adverse events, and quality of life (QoL) to be important outcomes. Some patient input was sought and patients identified that all of the outcomes mentioned would be important to them in making any treatment decisions.

Recommendation 1a
For patients with newly diagnosed, primary stage III epithelial ovarian, fallopian tube, or primary peritoneal carcinoma, HIPEC should be considered for those with at least stable disease following neoadjuvant chemotherapy at the time of interval CRS if complete or optimal cytoreduction is achieved.
<i>Qualifying Statements for Recommendation 1a</i>
The Working Group members recommend prospectively collecting data on these patients to evaluate real-world outcomes and applicability.
Recommendation 1b
There is insufficient evidence to recommend the addition of HIPEC when primary CRS is performed for patients with newly diagnosed, primary advanced epithelial ovarian, fallopian tube, or primary peritoneal carcinoma outside of a clinical trial.

Recommendation 2

There is insufficient evidence to recommend HIPEC with CRS in patients with recurrent ovarian cancer outside of a clinical trial.

Recommendation 3

There is insufficient evidence to recommend HIPEC with CRS in patients with peritoneal colorectal carcinomatosis outside of a clinical trial.

Recommendation 4

There is insufficient evidence to recommend HIPEC with CRS for the prevention of peritoneal carcinomatosis in CRC outside of a clinical trial; however HIPEC using oxaliplatin is not recommended.

Recommendation 5

There is insufficient evidence to recommend HIPEC with CRS for the treatment of gastric peritoneal carcinomatosis outside of a clinical trial.

Recommendation 6

There is insufficient evidence to recommend HIPEC with CRS for the prevention of gastric peritoneal carcinomatosis outside of a clinical trial.

Recommendation 7

There is insufficient evidence to recommend HIPEC with CRS in patients with malignant peritoneal mesothelioma as a standard of care; however, patients should be referred to HIPEC specialty centres for assessment for treatment as part of an ongoing research protocol.

Qualifying Statements for Recommendation 7

- The Working Group members recommend prospective research protocols with standardized treatment approaches at high-volume centres as this will provide survival benchmarks and feasibility data for future comparative studies.

Recommendation 8

There is insufficient evidence to recommend HIPEC with CRS in patients with disseminated mucinous neoplasm in the appendix as a standard of care; however, patients should be referred to HIPEC specialty centres for assessment for treatment as part of an ongoing research protocol.

Qualifying Statements for Recommendation 8

- The Working Group members recommend prospective research protocols with standardized treatment approaches at high-volume centres as this will provide survival benchmarks and feasibility data for future comparative studies.