


# Lung Cancer Diagnosis Pathway Map

Version 2019.05



## Target Population

Patients who present with signs or symptoms suspicious of lung cancer.

## Pathway Map Considerations

- Primary care providers play an important role in the cancer journey and should be informed of relevant tests and consultations. Ongoing care with a primary care provider is assumed to be part of the pathway. For patients who do not have a primary care provider, [Health Care Connect](#), is a government resource that helps patients find a doctor or nurse practitioner.
- Throughout the pathway, a shared decision-making model should be implemented to enable and encourage patients to play an active role in the management of their care. For more information see [Person-Centered Care Guideline](#) and [EBS #19-2 Provider-Patient Communication\\*](#)
- Hyperlinks are used throughout the pathway to provide information about relevant CCO tools, resources and guidance documents.
- The term 'health care provider', used throughout the pathway, includes primary care providers and specialists, nurse practitioners, and emergency physicians.
- For more information on the Diagnostic Assessment Program (DAP) refer to the [Organizational Standards for DAPs](#)
- Psychosocial oncology (PSO) is the interprofessional specialty concerned with understanding and treating the social, practical, psychological, emotional, spiritual and functional needs and quality-of-life impact that cancer has on patients and their families. Psychosocial care should be considered an integral and standardized part of cancer care for patients and their families at all stages of the illness trajectory. For more information, visit [EBS #19-3\\*](#)
- Counseling and treatment for smoking cessation should be initiated early on in the pathway and continued by care providers throughout the pathway as necessary [Program Training & Consultation Centre – Hospital Based Resources](#)










\* **Note.** [EBS #19-2](#) and [EBS #19-3](#) is older than 3 years and is currently listed as 'For Education and Information Purposes'. This means that the recommendations will no longer be maintained but may still be useful for academic or other information purposes.

## Pathway Map Legend



### Colour Guide

	Primary Care
	Palliative Care
	Pathology
	Diagnostic Assessment Program (DAP)
	Surgery
	Radiation Oncology
	Medical Oncology
	Radiology
	Multidisciplinary Cancer Conference (MCC)
	Respirologist
	Psychosocial Oncology (PSO)

### Shape Guide

	Intervention
	Decision or assessment point
	Patient (disease) characteristics
	Consultation with specialist
	Exit pathway
	Off-page reference
	Patient/Provider interaction
	Referral
	Wait time indicator time point

### Line Guide

	Required
	Possible

## Pathway Map Disclaimer

This pathway map is a resource that provides an overview of the treatment that an individual in the Ontario cancer system may receive.

The pathway map is intended to be used for informational purposes only. The pathway map is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. Further, all pathway maps are subject to clinical judgment and actual practice patterns may not follow the proposed steps set out in the pathway map. In the situation where the reader is not a healthcare provider, the reader should always consult a healthcare provider if he/she has any questions regarding the information set out in the pathway map. The information in the pathway map does not create a physician-patient relationship between Cancer Care Ontario (CCO) and the reader.

While care has been taken in the preparation of the information contained in the pathway map, such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability.


CCO and the pathway map's content providers (including the physicians who contributed to the information in the pathway map) shall have no liability, whether direct, indirect, consequential, contingent, special, or incidental, related to or arising from the information in the pathway map or its use thereof, whether based on breach of contract or tort (including negligence), and even if advised of the possibility thereof. Anyone using the information in the pathway map does so at his or her own risk, and by using such information, agrees to indemnify CCO and its content providers from any and all liability, loss, damages, costs and expenses (including legal fees and expenses) arising from such person's use of the information in the pathway map.

This pathway map may not reflect all the available scientific research and is not intended as an exhaustive resource. CCO and its content providers assume no responsibility for omissions or incomplete information in this pathway map. It is possible that other relevant scientific findings may have been reported since completion of this pathway map. This pathway map may be superseded by an updated pathway map on the same topic.

The pathway map is intended to be used for informational purposes only. The pathway map is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. Further, all pathway maps are subject to clinical judgment and actual practice patterns may not follow the proposed steps set out in the pathway map. In the situation where the reader is not a healthcare provider, the reader should always consult a healthcare provider if he/she has any questions regarding the information set out in the pathway map. The information in the pathway map does not create a physician-patient relationship between Cancer Care Ontario (CCO) and the reader.

Screen for psychosocial needs, and assessment and management of symptoms. [Click here for more information about symptom assessment and management tools](#)

Consider the introduction of palliative care, early and across the cancer journey [Click here for more information about palliative care](#)



<sup>1</sup> Refer to the American College of Chest Physicians Clinical Practice Guideline, *Chest*, 132, 149-160 for features of a standardized evaluation for systematic metastases and a list of paraneoplastic syndromes associated with lung cancer.

<sup>2</sup> The following factors have been shown to increase the risk of lung cancer: current or previous smoking of tobacco in/through cigarettes, cigars, dry pipe or water pipe (bong, vaping), second hand exposure to tobacco smoke, lung Diseases (chronic obstructive pulmonary disease, asthma, pulmonary fibrosis), previous exposure to asbestos or other known carcinogens, occupational exposure to: dust or other microscopic particles, diesel engine emissions, or chlorinated solvents, personal or family history of cancer (especially lung, head & neck), infections (HPV 16/18 of the respiratory tract, previous pneumonia, HIV), other illnesses or health issues (silicosis, tuberculosis, lupus, rheumatoid arthritis, systemic sclerosis, diabetes, periodontal disease, blood lipid levels, increased abdominal obesity), occupations (miners, painters, iron and steel workers, bricklayers, welders), environmental (in-home burning of coal and/or biomass, unventilated cooking over high heat, air pollution, low socioeconomic status, high caffeine intake)


<sup>3</sup> These patients should be accepted by the lung DAP if the lung DAP can facilitate a diagnosis within one week.

<sup>4</sup> An abnormal chest x-ray or an abnormal CT scan of chest suspicious of lung cancer is required with each DAP referral. A CT scan of the chest is not required for acceptance of a lung DAP referral if the chest x-ray is abnormal but a CT scan-chest is required prior to assessment at a lung DAP. Patient history should be mandatory as part of the referral and include, at a minimum: comorbidities, medications, allergies major health issues and symptoms that prompted the DAP referral.

The pathway map is intended to be used for informational purposes only. The pathway map is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. Further, all pathway maps are subject to clinical judgment and actual practice patterns may not follow the proposed steps set out in the pathway map. In the situation where the reader is not a healthcare provider, the reader should always consult a healthcare provider if he/she has any questions regarding the information set out in the pathway map. The information in the pathway map does not create a physician-patient relationship between Cancer Care Ontario (CCO) and the reader.

Screen for psychosocial needs, and assessment and management of symptoms. [Click here for more information about symptom assessment and management tools](#)

Consider the introduction of palliative care, early and across the cancer journey [Click here for more information about palliative care](#)




<sup>4</sup> An abnormal chest x-ray or an abnormal CT scan of chest suspicious of lung cancer is required with each DAP referral. A CT scan of the chest is not required for acceptance of a lung DAP referral if the chest x-ray is abnormal but a CT scan-chest is required prior to assessment at a lung DAP. Patient history should be mandatory as part of the referral and include, at a minimum: comorbidities, medications, allergies major health issues and symptoms that prompted the DAP referral.

The pathway map is intended to be used for informational purposes only. The pathway map is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. Further, all pathway maps are subject to clinical judgment and actual practice patterns may not follow the proposed steps set out in the pathway map. In the situation where the reader is not a healthcare provider, the reader should always consult a healthcare provider if he/she has any questions regarding the information set out in the pathway map. The information in the pathway map does not create a physician-patient relationship between Cancer Care Ontario (CCO) and the reader.

Screen for psychosocial needs, and assessment and management of symptoms. [Click here for more information about symptom assessment and management tools](#)

Consider the introduction of palliative care, early and across the cancer journey [Click here for more information about palliative care](#)



<sup>5</sup> Depending on local resources, radial miniprobe navigational bronchoscopy with lung biopsy may be considered.

<sup>6</sup> If the endobronchial ultrasound transbronchial needle aspiration is negative but there is a high level of suspicion of lung cancer, a mediastinoscopy should be completed.

<sup>7</sup> Results go to ordering and referring physician and family physician


<sup>8</sup> For more information about biomarkers, refer to the [Lung Cancer Tissue Pathway](#)

<sup>9</sup> Follow-up as per the Fleischner guidelines. For more information see Guidelines for Management of Small Pulmonary Nodules Detected on CT Scans: A Statement from the Fleischner Society. (2005). Radiology, 237, 395-400.

The pathway map is intended to be used for informational purposes only. The pathway map is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. Further, all pathway maps are subject to clinical judgment and actual practice patterns may not follow the proposed steps set out in the pathway map. In the situation where the reader is not a healthcare provider, the reader should always consult a healthcare provider if he/she has any questions regarding the information set out in the pathway map. The information in the pathway map does not create a physician-patient relationship between Cancer Care Ontario (CCO) and the reader.

Screen for psychosocial needs, and assessment and management of symptoms. [Click here for more information about symptom assessment and management tools](#)

Consider the introduction of palliative care, early and across the cancer journey [Click here for more information about palliative care](#)



<sup>7</sup> Results go to ordering and referring physician and family physician


<sup>8</sup> For more information about biomarkers, refer to the [Lung Cancer Tissue Pathway](#)



The pathway map is intended to be used for informational purposes only. The pathway map is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. Further, all pathway maps are subject to clinical judgment and actual practice patterns may not follow the proposed steps set out in the pathway map. In the situation where the reader is not a healthcare provider, the reader should always consult a healthcare provider if he/she has any questions regarding the information set out in the pathway map. The information in the pathway map does not create a physician-patient relationship between Cancer Care Ontario (CCO) and the reader.

Screen for psychosocial needs, and assessment and management of symptoms. [Click here for more information about symptom assessment and management tools](#)

Consider the introduction of palliative care, early and across the cancer journey [Click here for more information about palliative care](#)



<sup>10</sup> If emergency situation, symptomatic brain metastases, superior vena cava obstruction, spinal compression or stage I-III disease.