



## Evidence-Based Series 12-11 REQUIRES UPDATING

A Quality Initiative of the  
Program in Evidence-Based Care (PEBC), Cancer Care Ontario (CCO)

### Patient Safety Issues: Key Components of Chemotherapy Labelling

*M. Trudeau, E. Green, R. Cosby, F. Charbonneau, T. Easty,  
Y. Ko, P. Marchand, D.U. N. Berger, and S. Hertz*

Report Date: August 6, 2009

An assessment conducted in December 2018 indicated that Evidence-based Series (EBS) 12-11 requires updating. It is still appropriate for this document to be available while this updating process unfolds. The PEBC has a formal and standardized process to ensure the currency of each document ([PEBC Assessment & Review Protocol](#))

EBS 12-11 is comprised of 3 sections. You can access the summary and full report here:

<https://www.cancercareontario.ca/en/guidelines-advice/types-of-cancer/1191>

Section 1: Recommendations

Section 2: Evidentiary Base

Section 3: EBS Development Methods and External Review Process

For information about the PEBC and the most current version of all reports, please visit the CCO website at <http://www.cancercare.on.ca/>

or contact the PEBC office at:

Phone: 905-527-4322 ext. 42822 Fax: 905-526-6775 E-mail: [ccopgi@mcmaster.ca](mailto:ccopgi@mcmaster.ca)

**PEBC Report Citation (Vancouver Style):** Trudeau M, Green E, Cosby R, Charbonneau F, Easty T, Ko Y, et al. Patient safety issues: key components of chemotherapy labelling. Toronto (ON): Cancer Care Ontario; 2009 Aug 6. Program in Evidence-based Care Practice Guideline Report No.: 12-11 REQUIRES UPDATING.

**Journal Citation (Vancouver Style):** Trudeau M, Green E, Cosby R, Charbonneau F, Easty T, Ko Y, et al. Key components of intravenous chemotherapy labeling: a systematic review and practice guideline. J Oncol Pharm Pract. 2011 Dec;17(4):409-24.

**Evidence-Based Series 12-11: Section 1**

**Patient Safety Issues: Key Components of Chemotherapy Labelling:  
Guideline Recommendations**

*M. Trudeau, E. Green, R. Cosby, F. Charbonneau, T. Easty,  
Y. Ko, P. Marchand, D.U. N. Berger, and S. Hertz*

A Quality Initiative of the Chemotherapy Labelling Panel  
Program in Evidence-Based Care (PEBC), Cancer Care Ontario (CCO)

**Report Date: August 6, 2009**

**QUESTION**

What are the necessary components and formatting of a chemotherapy label to maximize safe delivery and minimize errors? Chemotherapy labels associated with the delivery of a dose of intravenous chemotherapy are of particular interest.

**INTENDED USERS**

The intended users of this guidance document are any health care professionals who prescribe, prepare, or administer intravenous chemotherapy, including medical oncologists, pharmacists, pharmacy technicians, and oncology nurses, as well as designers of prescription label software, patient safety directors in organizations, administrators of hospitals, and community access care organizations.

**RECOMMENDATIONS**

The following recommendations are based on the expert opinion of the Chemotherapy Labelling Panel but informed by the currently available evidence (see Section 2). The evidentiary base is composed of three guidelines developed by expert groups, one systematic review, and 13 studies of varying design and sample size. These recommendations apply to the production of intravenous chemotherapy labels in a cancer setting. Although the production of labels for investigational cancer drugs was not specifically examined, the same principles apply for all intravenous chemotherapy labels. Examples of labels using these recommendations are included at the end of this section.

## 1. General Components for Medication Labels

The following are general components of an optimal drug label for injectable dosage forms.

### **(a) Identifying Information**

- Patient's name (first name, middle name or initial, and last name **OR** last name, first name, and middle name or initial such that it is consistent with the rest of the patient record) and unique identifier
- Drug name
- Amount of drug per container
- In those circumstances in which overfill is required, the overfill volume (in mL) should be printed on the label separately from the dose information
- If a product contains two or more active ingredients, they should all appear in the generic name field

### **(b) Drug Information**

- Route of administration
- Amount of drug per dose (when the container holds more than one dose, e.g., multiple doses administered intermittently over a 24-hour time period)

### **(c) Administration Information**

- Volume of fluid to be administered
- Duration of infusion
- Rate of administration expressed in mL/hour or as a duration in minutes in the case of medications given by IV push. There is a need to standardize pump technology within an institution or at least to use pumps with a common format. The use of pumps programmed in mL/hour is strongly recommended over the use of pumps programmed in mL/24 hour.
- Supplemental administration instructions (e.g., starting and completion dates/times, prohibitions about when medications are to be administered with respect to other medications, warnings about route of administration, handling and storage conditions)
- Numbering of the medication containers, when the drug is to be administered sequentially (e.g., bag 1 of 3)
- Relevant auxiliary information should be included on auxiliary labels. Examples of auxiliary labels include "AVOID EXTRAVASATION" and "FOR INTRAVENOUS USE ONLY - FATAL IF GIVEN BY OTHER ROUTES"

### **(d) General Formatting**

- Allow for text wrap and continuation of information on another label. This is intended to allow for long names and enough space to ensure readability as well as eliminating the need to add in additional hand-written information.
- Use white labels: better visualization of text and bar codes (if used). Use black for bar codes.
- If a different colour label is required to draw attention to a specific class of high-alert drug, use yellow labels.

## 2. General Principles for Label Preparation

The following are general formatting principles to be considered when preparing a chemotherapy drug label for injectable dosage forms.

### (a) Drug Name

The following practices are recommended:

- Use the complete generic drug name rather than an abbreviated version.
  - cisplatin not CDDP
- Use lower case or mixed case lettering for generic drug names as appropriate
  - Use TALL man lettering to differentiate between look alike/sound alike drug names (examples can be found at <http://www.ismp.org/tools/tallmanletters.pdf>)
    - CISplatin to differentiate it from CARBOplatin
- List the brand name using uppercase letters.
  - HERCEPTIN

### (b) Abbreviations and Dose Designations

- The recommended practice is to follow Institute for Safe Medication Practices (ISMP) guidelines for abbreviations and dose expressions (examples are provided in Section 2, Table 6) and United States Pharmacopeia (USP) standards for dosage units and standard units for weight and measures (examples are provided in Section 2, Table 7). Alternative abbreviations and dose expressions should be avoided.

### (c) Font, Font Size, and Formatting

It is recommended that:

- Patient name, generic drug name and patient specific dose are bolded.
- 12-point Arial, Verdana or an equivalent proportionally spaced font is used for all text and numbers.
  - **Jane A. Smith** not Jane A. Smith
- When drug name, strength, dosage form, and dosage units appear together, provide a space between them
  - propranolol 20 mg *not* propranolol20 mg
- Laser printers that support all label formatting expectations be used.

### (d) Order of Information

- It is recommended that label information should be presented in the following order: generic name, brand name, patient dose, dosage units, and route of administration.
  - **ondansetron (ZOFTRAN) 4 mg IV Push**  
Dose = 4 mg = 2 mL  
(2mg per mL)\*
- \*include this information only if needed by practitioners (e.g., to program infusion pump)
- The order of information on the label should match the user's workflow; that is the order in which information is programmed into the pump. This will vary depending on the type of pump used in an institution.

### (e) Technology

- While more evidence is required, the use of bar coding may be considered for use.
- The use of computerized physician order entry (CPOE) is recommended.

## KEY EVIDENCE

- Guideline documents (1-3) provided a framework to identify domains that ought to be considered in an optimal label.
- Label generation should be guided by the overarching rule that medication labels not contain any unnecessary information (4).
- Communication of orders for infusions should be standardized such that “mL per hour” is used rather than “mL per 24 hour” (4).
- ISMP Canada (5) and ISMP United States [US] (6) provide sets of abbreviations, symbols and dose designations that should not be used, which the authors of this document endorse. Please see Tables 6 and 7 in Section 2 for examples.
- TALL man lettering has consistently been shown to reduce drug name identification errors (7-10).
- Larger font size and font weight results in fewer reading errors (11) and better knowledge acquisition (12).
- Proportionally spaced fonts result in better reading speed and accuracy (11).
- There are beginning studies on bar coding indicating that medication administration errors may be reduced with the use of this technology (13, 14). More research is needed before a recommendation regarding this technology can be made.
- CPOE has been demonstrated to reduce medication errors (15-19).
- There is limited evidence that laser printers are preferred over dot-matrix printers (20).

## Examples of Labels using the Recommendations in this Guidance Document

The following examples are for illustrative purposes and do not account for overfill volumes which may require consideration.

### Example 1 - Intravenous Infusion

<b>Smith, John A.</b>	20000133	→ Patient Name/Unique Identifier
<b>irinotecan hcl 320 mg/16 mL IV</b>		→ Drug Name/Amount of Drug/Route of Administration
(20 mg/mL)		→ Drug Concentration (only if needed)
solution: D5W	volume: 500 mL	→ Diluent/Amount of Diluent
total volume: 516 mL		→ Volume of Fluid to be Administered
rate: 344 mL/hour		→ Rate of Administration
Infuse IV over 90 minutes; run concurrently with leucovorin calcium.		→ Administration Instructions
date:12-Jun-2009		→ Date

## Example 2 - Intravenous Infusion

**Smith, John A.** 20000133

**leucovorin calcium 360 mg/36 mL IV**  
(10 mg/mL)

solution: D5W      volume: 250 mL

total volume: 286 mL

rate: 191 mL/hour

Infuse IV over 90 minutes; run concurrently with  
irinotecan.

date:12-Jun-2009

## Example 3 - Continuous Intravenous Infusion

**Smith, John A.** 20000133

**fluorouracil 4350 mg/87 mL CIV**  
(50 mg/mL)

solution: D5W      volume: 146 mL

total volume: 233 mL

rate: 5 mL/hour

IV continuous infusion over 46 hours.

\*\*\* INSERT INFUSOR REFERENCE NUMBER \*\*\*

date:12-Jun-2009

#### Example 4 - Intravenous Push with Multiple Syringe and use of TALL man Lettering

<b>Smith, Mary A.</b>	20000298
<b>EPIrubicin 166 mg/83 mL IV</b> (2 mg/mL)	
1 of 2 syringes.	
Each syringe contains 83 mg/41.5 mL.	
Infuse slowly IV at a rate of 5 mL/minute.	
<b>AVOID EXTRAVASATION</b>	→ auxiliary label
date:12-Jun-2009	

#### Example 5 - Multiple Additives

<b>Smith, John A.</b>	20000133
<b>calcium gluconate 1 g/10 mL IV</b> (0.1 g/mL)	
<b>magnesium sulfate 1 g/2 mL IV</b> (0.5 g/mL)	
solution: D5W	volume: 250 mL
total volume: 262 mL	
rate: 786 mL/hour	
Infuse over 20 minutes prior to oxaliplatin.	
date:12-Jun-2009	

## FUTURE RESEARCH

More research is needed on the use and effectiveness of strategies to reduce medication administration errors. Specifically, more studies evaluating the effectiveness of bar coding to reduce medication errors and adverse events are needed. In addition, studies are needed to evaluate the best method(s) for patient identification to enhance the safe administration of chemotherapy. There are now a few institutions that generate two labels: one for pharmacy staff who fill the prescriptions and one for the nurses who administer the chemotherapy. Research is needed to determine if a system that makes use of two labels results in fewer medication errors than a system in which one label is used. The safe administration of chemotherapy is a complex process in which good labels are necessary but not a sole or sufficient strategy.

### *Funding*

The PEBC is a provincial initiative of Cancer Care Ontario supported by the Ontario Ministry of Health and Long-Term Care through Cancer Care Ontario. All work produced by the PEBC is editorially independent from its funding source.

### *Copyright*

This report is copyrighted by Cancer Care Ontario; the report and the illustrations herein may not be reproduced without the express written permission of Cancer Care Ontario. Cancer Care Ontario reserves the right at any time, and at its sole discretion, to change or revoke this authorization.

### *Disclaimer*

Care has been taken in the preparation of the information contained in this report. Nonetheless, any person seeking to apply or consult the report is expected to use independent medical judgement in the context of individual clinical circumstances or seek out the supervision of a qualified clinician. Cancer Care Ontario makes no representation or guarantees of any kind whatsoever regarding the report content or use or application and disclaims any responsibility for its application or use in any way.

### *Contact Information*

For further information about this report, please contact:

**Dr. Maureen Trudeau**, Cancer Care Ontario, 620 University Avenue, Toronto, ON, M5G 2L7  
Phone: 416-480-5145 Fax: 416-481-6002 E-mail: [Maureen.trudeau@sunnybrook.ca](mailto:Maureen.trudeau@sunnybrook.ca)

or

**Esther Green**, Cancer Care Ontario, 620 University Avenue, Toronto, ON, M5G 2L7  
Phone: 416-971-9800 x1278 E-mail: [esther.green@cancercare.on.ca](mailto:esther.green@cancercare.on.ca)

For information about the PEBC and the most current version of all reports,  
please visit the CCO website at <http://www.cancercare.on.ca/>  
or contact the PEBC office at:

Phone: 905-527-4322 ext. 42822 Fax: 905-526-6775 E-mail: [ccopgi@mcmaster.ca](mailto:ccopgi@mcmaster.ca)



## REFERENCES

1. Kohler DR, Montello MJ, Green L, Huntley C, High JL, Fallavollita A, Jr, et al. Standardizing the expression and nomenclature of cancer treatment regimens. *Am J Health Syst Pharm.* 1998;55(2):137-44.
2. ASHP Council on Professional Affairs. ASHP guidelines on preventing medication errors with antineoplastic agents. *Am J Health Syst Pharm.* 2002;59(17):1648-68.
3. Institute for Safe Medication Practices US. Principles of designing a medication label for injectable syringes for patient specific, inpatient use. Horsham (PA): Institute for Safe Medication Practices US; 2008 [cited: 2008 Mar 5]. Available from: <http://www.ismp.org/Tools/guidelines/labelFormats/injectionSyringe.asp>
4. Institute for Safe Medication Practices Canada. Fluorouracil incident root cause analysis. Toronto: Institute for Safe Medication Practices Canada; 2007 [cited: 2008 Mar 5]. Available from: <http://www.cancerboard.ab.ca/NR/rdonlyres/2FB61BC4-70CA-4E58-BDE1-1E54797BA47D/0/FluorouracilIncidentMay2007.pdf>
5. Institute for Safe Medication Practices Canada. Do not use: dangerous abbreviations, symbols and dose designations. Toronto: Institute for Safe Medication Practices Canada; 2006 [cited: 2008 Mar 5]. Available from: <http://www.ismp-canada.org/download/ISMPCanadaListOfDangerousAbbreviations.pdf>
6. Institute for Safe Medication Practices US. ISMP's list of error-prone abbreviations, symbols and dose designations. Horsham (PA): Institute for Safe Medication Practices US; 2006 [cited: 2008 Mar 5]. Available from: <http://www.ismp.org/Tools/errorproneabbreviations.pdf>
7. Filik R, Purdy K, Gale A, Gerrett D. Investigating medication errors caused by confusable drug names. In: Schmalhofer F, Young RM, Katz G, editors. *Proceedings of EuroCogSci 03.* London: Lawrence Erlbaum Associates; 2003. p. 383.
8. Filik R, Gale A, Purdy K, Gerrett D. Medication errors and human factors. In: McCabe P, editor. *Contemporary ergonomics.* London: Taylor & Francis; 2003. p. 471-6.
9. Filik R, Purdy K, Gale A, Gerrett D. Drug name confusion: evaluating the effectiveness of capital ("Tall Man") letters using eye movement data. *Soc Sci Med.* 2004;59(12):2597-601.
10. Filik R, Purdy K, Gale A, Gerrett D. Labeling of medicines and patient safety: evaluating methods of reducing drug name confusion. *Hum Factors.* 2006;48(1):39-47.
11. Smither JA, Braun CC. Readability of prescription drug labels by older and younger adults. *J Clin Psychol Med Settings.* 1994;1(2):149-59.
12. Wogalter MS, Vigilante WJ, Jr. Effects of label format on knowledge acquisition and perceived readability by younger and older adults. *Ergonomics.* 2003;46(4):327-44.
13. Poon EG, Cina JL, Churchill W, Patel N, Featherstone E, Rothschild JM, et al. Medication dispensing errors and potential adverse drug events before and after implementing bar code technology in the pharmacy. *Ann Intern Med.* 2006;145(6):426-34.
14. Paoletti RD, Suess TM, Lesko MG, Feroli AA, Kennel JA, Mahler JM, et al. Using bar-code technology and medication observation methodology for safer medication administration. *Am J Health Syst Pharm.* 2007;64(5):536-43.
15. Kaushal R, Shojania KG, Bates DW. Effects of computerized physician order entry and clinical decision support systems on medication safety: a systematic review. *Arch Intern Med.* 2003;163(12):1409-16.
16. Kim GR, Chen AR, Arceci RJ, Mitchell SH, Kokoszka KM, Daniel D, et al. Error reduction in pediatric chemotherapy: computerized order entry and failure modes and effects analysis [see comment]. *Arch Pediatr Adolesc Med.* 2006;160(5):495-8.

17. Huertas Fernandez MJ, Baena-Canada JM, Martinez Bautista MJ, Arriola AE, Garcia Palacios MV. Impact of computerised chemotherapy prescriptions on the prevention of medication errors. *Clin Transl Onc.* 2006;8(11):821-5.
18. Shamliyan TA, Duval S, Du J, Kane RL. Just what ever doctor ordered. Review of the evidence of the impact of computerized physician order entry system on medication errors. *Health Serv Res.* 2008;43(1):32-53.
19. Ammenwerth E, Schnell-Inderst P, Machan C, Siebert U. The effect of electronic prescribing on medication errors and adverse drug events: A systematic review. *J Am Med Inform Assoc.* 2008;15(5):585-600.
20. Luscombe DK, Jinks MJ, Duncan S. A survey of prescription label preferences among community pharmacy patrons. *J Clin Pharm Ther.* 1992;17(4):241-4.

IN REVIEW