

CRBPGEMC Treatment

This handout gives general information about this cancer treatment.

You will learn:

- who to contact for help
- what the treatment is
- how it is given
- what to expect while on treatment



This handout was created by Ontario Health (Cancer Care Ontario) together with patients and their caregivers who have also gone through cancer treatment. It is meant to help support you through your cancer treatment and answer some of your questions.

This information does not replace the advice of your health care team. Always talk to your health care team about your treatment.

Who do I contact if I have questions or need help?

My cancer health care provider is: _____

During the day I should contact: _____

Evenings, weekends and holidays: _____

What is this treatment?

CRBPGEMC is the code name of your cancer treatment regimen. CRBPGEMC is used to treat different types of cancers, including lung, bladder, ovarian, head and neck, biliary tract or other cancers.

A regimen is a combination of medications to treat cancer.

This regimen name is made up of letters from the names of the 2 medications in your treatment.


Here are the names of the medications in this regimen:

CRBP = CaRBoPlatin
GEMC = GEMCitabine

CRBPGEMC may be given on a schedule that is different from what is shown on the calendar below. Speak with your health care team to see if the below schedule applies to you.

The treatment is usually divided into **cycles**. Each cycle is **3 weeks** long. Your health care team will tell you how many cycles you need.

Here is a picture of the schedule for CRBPGEMC treatment:



During each 3-week cycle, you will have CRBPGEMC treatment on day 1 at the hospital, followed by GEMC treatment on day 8 at the hospital. You will then have a week with no treatment at the hospital (2 weeks of treatment, then 1 week with no treatment).

Each cycle looks like this:

Day 1	2	3	4	5	6	7
Treatment Day: Go to the hospital for CRBPGEMC Treatment.	NO Treatment					
8	9	10	11	12	13	14
Treatment Day: Go to the hospital for GEMC Treatment.	NO Treatment					
15	16	17	18	19	20	21
						



Remember To:

- ✓ Tell your health care team about all of the other medications you are taking.
- ✓ Keep taking other medications that have been prescribed for you, unless you have been told not to by your health care team.

You will have a blood test to check for hepatitis B before starting treatment. See the [Hepatitis B and Cancer Medications](#) pamphlet for more information.

How is this treatment given?

The medications in your treatment are given through an IV (injected into a vein) at the hospital.

You will have a blood test before each treatment cycle to make sure it is safe for you to get treatment.

What other medications are given with this treatment?

To Prevent Nausea and Vomiting

You will be given medications to help prevent nausea (feeling like throwing up) and vomiting (throwing up) before they start.

- ✓ These are called anti-nausea medications and include medications such as ondansetron (Zofran®), granisetron (Kytril®), aprepitant (Emend®), dexamethasone or others.

What other important things should I know about this treatment?

If you get radiation treatment at the same time as CRBPGEMC, you may get worse side effects from radiation. Tell your health care team if you have had or will be getting radiation therapy.

DO this while on treatment

- ✓ DO tell your health care team about any other medical conditions that you have such as, hearing, heart, lung, liver or kidney problems, issues with your bone marrow or any allergies.
- ✓ DO check with your health care team before getting any vaccinations, surgery, dental work or other medical procedures or if you have had previous radiation therapy.
- ✓ DO talk to your health care team about your risk of getting other cancers or heart problems after this treatment.
- ✓ DO drink plenty of fluids to prevent kidney problems. Drink at least 6 to 8 cups (2 Litres) of water or other liquids per day on your CRBPGEMC treatment day and for 1-2 days afterwards, unless your healthcare team has told you to drink more or less.
- ✓ DO consider asking someone to drive you to and from the hospital on your treatment days. You may feel drowsy or dizzy after your treatment.

DO NOT do this while on treatment



- ✗ DO NOT take any other medications, such as vitamins, over-the-counter (non-prescription) drugs, or natural health products without checking with your health care team.
- ✗ DO NOT start any complementary or alternative therapies, such as acupuncture or homeopathic medications, without checking with your health care team.
- ✗ DO NOT smoke or drink alcohol while on treatment without talking to your health care team first. Smoking and drinking can make side effects worse and make your treatment not work as well.

Will this treatment interact with other medications or natural health products?

Yes, the medications in this regimen can interact with other medications, vitamins, foods and natural health products. Interactions can make the treatment not work as well or cause severe side effects.

Tell your health care team about all of your:

- prescription and over-the-counter (non-prescription) medications
- natural health products such as vitamins, herbal teas, homeopathic medicines, and other supplements

Check with your health care team before starting or stopping any of them.

If you take seizure medications (such as phenytoin), your health care team may monitor your blood levels closely and may change your dose.

If you are taking a blood thinner (such as warfarin), your health care team may need extra blood tests and may change your dose.



Talk to your health care team **BEFORE** taking or using these :

- Anti-inflammatory medications such as ibuprofen (Advil[®] or Motrin[®]), naproxen (Aleve[®]) or Aspirin[®].
- Over-the-counter products such as dimenhydrinate (Gravol[®])
- Natural health products such as St. John's Wort
- Supplements such as vitamin C
- Grapefruit juice
- Alcoholic drinks
- Tobacco
- All other drugs, such as marijuana or cannabis (medical or recreational)

What should I do if I feel unwell, have pain, a headache or a fever?

- ✓ **Always** check your temperature to see if you have a fever **before** taking any medications for fever or pain (such as acetaminophen (Tylenol®) or ibuprofen (Advil®)).
 - Fever can be a sign of infection that may need treatment right away.
 - If you take these medications before you check for fever, they may lower your temperature and you may not know you have an infection.

How to check for fever:

Keep a digital (electronic) thermometer at home and take your temperature if you feel hot or unwell (for example, chills, headache, mild pain).

- You have a fever if your temperature taken **in your mouth (oral temperature)** is:
 - 38.3°C (100.9°F) or higher at any time
- OR
- 38.0°C (100.4°F) or higher for at least one hour.



If you do have a fever:

- ✓ **Try to contact your health care team. If you are not able to talk to them for advice, you MUST get emergency medical help right away.**
- ✓ Ask your health care team for the [Fever](#) pamphlet for more information.

If you do not have a fever but have mild symptoms such as headache or mild pain:

- ✓ Ask your health care team about the right medication for you. **Acetaminophen (Tylenol®)** is a safe choice for most people.



Talk to your health care team before you start taking ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or ASA (Aspirin®), as they may increase your chance of bleeding or interact with your cancer treatment.



Talk to your health care team if you already take **low dose aspirin** for a medical condition (such as a heart problem). It may still be safe to take.

How will this treatment affect sex, pregnancy and breast feeding?

Talk to your health care team about:

- How this treatment may affect your sexual health.
- How this treatment may affect your ability to have a baby, if this applies to you.

This treatment may harm an unborn baby. Tell your health care team if you or your partner are pregnant, become pregnant during treatment, or are breastfeeding.

- If there is **any** chance of pregnancy happening, you and your partner together must use **2 effective forms of birth control** at the same time while you are on treatment. Talk to your health care team about which birth control options are best for you, and how long you should use them after your last treatment dose.
- Do not use hormonal birth control (such as birth control pills), unless your health care team told you that they are safe. Talk to your healthcare team about the safest birth control for you.
- Do not breastfeed while on this treatment. Talk to your health care team about how long to wait before you start breastfeeding after your last treatment dose, if this applies to you.

What are the side effects of this treatment?

The following table lists side effects that you may have when getting CRBPGEMC treatment. The table is set up to list the most common side effects first and the least common last. It is unlikely that you will have all of the side effects listed and you may have some that are not listed.

Read over the side effect table so that you know what to look for and when to get help. Refer to this table if you experience any side effects while on CRBPGEMC treatment.

Very Common Side Effects (50 or more out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Liver problems</p> <p>(May be severe)</p> <p>Your health care team may check your liver function with a blood test. Liver changes do not usually cause any symptoms.</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Rarely, you may develop yellowish skin or eyes, unusually dark pee or pain on the right side of your belly. This may be severe. <p>What to do?</p> <p>If you have any symptoms of liver problems, get emergency medical help right away.</p>	<p>Get emergency medical help right away</p>
<p>Low neutrophils (white blood cells) in the blood (neutropenia)</p> <p>(May be severe)</p> <p>When neutrophils are low, you are at risk of getting an infection more easily. Ask your health care team for the Neutropenia (Low white blood cell count) pamphlet for more information.</p> <p>What to look for?</p> <ul style="list-style-type: none"> • If you feel hot or unwell (for example if you have chills or a new cough), you must check your temperature to see if you have a fever. • Do not take medications that treat a fever before you take your temperature (for example, Tylenol® (acetaminophen), or Advil®) 	<p>If you have a fever, try to contact your health care team. If you are unable to talk to the team for advice, you MUST get emergency medical help right away.</p>

Very Common Side Effects (50 or more out of 100 people)	
Side effects and what to do	When to contact health care team
<p>(ibuprofen)).</p> <ul style="list-style-type: none"> Do not eat or drink anything hot or cold right before taking your temperature. <p>You have a fever if your temperature taken in your mouth (oral temperature) is:</p> <ul style="list-style-type: none"> 38.3°C (100.9°F) or higher at any time OR 38.0°C (100.4°F) or higher for at least one hour. <p>What to do?</p> <p>If your health care team has told you that you have low neutrophils:</p> <ul style="list-style-type: none"> Wash your hands often to prevent infection. Check with your health care team before getting any vaccines, surgeries, medical procedures or visiting your dentist. Keep a digital thermometer at home so you can easily check for a fever. <p>If you have a fever:</p> <p>If you have a fever, try to contact your health care team. If you are unable to talk to the team for advice, you must get emergency medical help right away.</p>	
<p>Low platelets in the blood</p> <p>(May be severe)</p> <p>When your platelets are low, you are at risk for bleeding and bruising. Ask your health care team for the Low Platelet Count pamphlet for more information.</p> <p>What to look for?</p> <ul style="list-style-type: none"> Watch for signs of bleeding: <ul style="list-style-type: none"> bleeding from your gums unusual or heavy nosebleeds bruising easily or more than normal black coloured stools (poo) or blood in your stools (poo) coughing up red or brown coloured mucus dizziness, constant headache or changes in your vision heavy vaginal bleeding red or pink coloured urine (pee) 	<p>Talk to your health care team if you have any signs of bleeding. If you have bleeding that doesn't stop or is severe (very heavy), you MUST get emergency help right away.</p>

Very Common Side Effects (50 or more out of 100 people)	
Side effects and what to do	When to contact health care team
<p>What to do?</p> <p>If your health care team has told you that you have low platelets:</p> <ul style="list-style-type: none"> • Tell your pharmacist that your platelet count may be low before taking any prescriptions or over-the-counter medication. • Check with your healthcare team before you go to the dentist. • Take care of your mouth and use a soft toothbrush. • Try to prevent cuts and bruises. • Ask your health care team what activities are safe for you. • Your treatment may have to be delayed if you have low platelets. Your health care team may recommend a blood transfusion. <p>If you have signs of bleeding:</p> <ul style="list-style-type: none"> • If you have a small bleed, clean the area with soap and water or a saline (saltwater) rinse. Apply pressure for at least 10 minutes. <p>If you have bleeding that does not stop or is severe (very heavy), you must get emergency medical help right away.</p>	
<p>Nausea and vomiting</p> <p>(More likely with carboplatin)</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Nausea is feeling like you need to throw up. You may also feel light-headed. • You may feel nausea within hours to days after your treatment. <p>What to do?</p> <p>To help prevent nausea:</p> <ul style="list-style-type: none"> • It is easier to prevent nausea than to treat it once it happens. • If you were given anti-nausea medication(s), take them as prescribed, even if you do not feel like throwing up. • Drink clear liquids and have small meals. Get fresh air and rest. • Do not eat spicy, fried foods or foods with a strong smell. • Limit caffeine (like coffee, tea) and avoid alcohol. 	<p>Talk to your healthcare team if nausea lasts more than 48 hours or vomiting lasts more than 24 hours or if it is severe</p>

Very Common Side Effects (50 or more out of 100 people)	
Side effects and what to do	When to contact health care team
<p>If you have nausea or vomiting:</p> <ul style="list-style-type: none"> • Take your rescue (as-needed) anti-nausea medication(s) as prescribed. • Ask your health care team for the Nausea & Vomiting pamphlet for more information. • Talk to your health care team if: <ul style="list-style-type: none"> ◦ nausea lasts more than 48 hours ◦ vomiting lasts more than 24 hours or if it is severe 	

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Too much or too little salt in your body (May be severe; with carboplatin)</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Muscle spasms, cramping, weakness, twitching, or convulsions. • Irregular heartbeat, confusion or blood pressure changes. <p>What to do?</p> <p>Get emergency medical help right away for severe symptoms.</p>	<p>Get emergency medical help right away for severe symptoms</p>
<p>Fatigue</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Feeling of tiredness or low energy that lasts a long time and does not go away with rest or sleep. <p>What to do?</p> <ul style="list-style-type: none"> • Be active. Aim to get 30 minutes of moderate exercise (you are able to talk comfortably while exercising) on most days. • Check with your health care team before starting any new exercise. • Pace yourself, do not rush. Put off less important activities. Rest when you 	<p>Talk to your health care team if it does not improve or if it is severe</p>

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>need to.</p> <ul style="list-style-type: none"> • Ask family or friends to help you with things like housework, shopping, and child or pet care. • Eat well and drink at least 6 to 8 glasses of water or other liquids every day (unless your health care team has told you to drink more or less). • Avoid driving or using machinery if you are feeling tired. <p>Ask your health care team for the Fatigue pamphlet for more information</p>	
<p>Flu-like symptoms</p> <p>You may feel like you have the flu for around 2 days after your IV treatment. These flu-like symptoms may not be signs of an infection.</p> <p>What to look for?</p> <ul style="list-style-type: none"> • You may have chills, headache and muscle pain. • You may feel tired and have a poor appetite. • Symptoms may happen at any time after you receive your treatment and usually go away as your body gets used to the medication. <p>What to do?</p> <ul style="list-style-type: none"> • Check your temperature to see if you have a fever. Read the above section "What should I do if I feel unwell, have pain, a headache or a fever?" • If you do have a fever, try to speak to your health care team. If you are unable to talk to them for advice, you MUST get emergency medical help right away. 	<p>Talk to your health care team if it does not improve or if it is severe</p>
<p>Rash; dry, itchy skin</p> <p>What to look for?</p> <ul style="list-style-type: none"> • You may have cracked, rough, flaking or peeling areas of the skin. • Your skin may look red and feel warm, like a sunburn. • Your skin may itch, burn, sting or feel very tender when touched. • The rash may be seen in areas where you have had radiation before. <p>What to do?</p> <p>To prevent and treat dry skin:</p> <ul style="list-style-type: none"> • Use fragrance-free skin moisturizer. • Protect your skin from the sun and the cold. 	<p>Talk to your health care team if it does not improve or if it is severe</p>

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<ul style="list-style-type: none"> • Use sunscreen with UVA and UVB protection and a SPF of at least 30. • Avoid perfumed products and lotions that contain alcohol. • Drink 6 to 8 cups of non-alcoholic, non-caffeinated liquids each day, unless your health care team has told you to drink more or less. <p>Rash may be severe in some rare cases and cause your skin to blister or peel. If this happens, get emergency medical help right away</p>	
<p>Kidney problems</p> <p>Your health care team may check for proteins in your urine (pee) and your kidney function regularly with a blood test. You may have blood in your urine.</p> <p>What to look for?</p> <ul style="list-style-type: none"> • New swelling in your hands, ankles, feet or other areas of your body. • Weight gain that is not normal for you. • Pain in your lower back. • Muscle twitches and cramps or itchiness that won't go away. • Nausea (feeling like you need to throw up) and vomiting. • Changes in urination (peeing) such as less urine than usual. • Foamy, frothy or bubbly-looking pee. <p>What to do?</p> <ul style="list-style-type: none"> • If you have any of these signs, talk to your health care team or go to your closest emergency department. 	Get emergency medical help right away

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Mild swelling</p> <p>What to look for?</p> <ul style="list-style-type: none"> • You may have mild swelling or puffiness in your arms and/or legs. Rarely, this may be severe. 	Talk to your health care team if it does not improve or if it is severe

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>What to do?</p> <p>To help prevent swelling:</p> <ul style="list-style-type: none"> • Eat a low-salt diet. <p>If you have swelling:</p> <ul style="list-style-type: none"> • Wear loose-fitting clothing. • For swollen legs or feet, keep your feet up when sitting. 	
<p>Mild joint, muscle pain or cramps</p> <p>What to look for?</p> <ul style="list-style-type: none"> • New pain in your muscles or joints, muscle cramps, or feeling achy. <p>What to do?</p> <ul style="list-style-type: none"> • Take pain medication (acetaminophen or opioids such as codeine, morphine, hydromorphone, oxycodone) as prescribed. • Read the above section: "What should I do if I feel unwell, have pain, a headache or a fever?" before taking acetaminophen (Tylenol®), ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or Aspirin. These medications may hide an infection that needs treatment or they may increase your risk of bleeding. • Rest often and try light exercise (such as walking) as it may help. <p>Ask your health care team for the Pain pamphlet for more information.</p>	<p>Talk to your health care team if it does not improve or if it is severe</p>
<p>Change in your hearing</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Not being able to hear as well as before. • New noise or ringing sounds in your ears. • Changes in hearing usually go away over time. In some rare cases they may be permanent. <p>What to do?</p> <ul style="list-style-type: none"> • Tell your health care team if you have any of these symptoms. Your health care team may need to change your medication. 	<p>Contact your health care team as soon as possible</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Hair thinning or loss</p> <p>(Generally mild)</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Your hair may become thin or fall out during or after treatment. • In most cases, your hair will grow back after treatment. The texture or colour may change. • In very rare cases, hair loss may be permanent. <p>What to do?</p> <ul style="list-style-type: none"> • Use a gentle soft brush. • Do not use hair sprays, bleaches, dyes and perms. 	<p>Talk to your health care team if this bothers you</p>
<p>Diarrhea</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Loose, watery, unformed stool (poo) that may happen days to weeks after you get your treatment. <p>What to do?</p> <p>If you have diarrhea:</p> <ul style="list-style-type: none"> • Take anti-diarrhea medication if your health care team prescribed it or told you to take it. • Do not eat foods or drinks with artificial sweetener (like chewing gum or 'diet' drinks), coffee and alcohol. • Eat many small meals and snacks instead of 2 or 3 large meals. • Drink at least 6 to 8 cups of liquids each day, unless your health care team has told you to drink more or less. • Talk to your health care team if you can't drink 6 to 8 cups of liquids each day when you have diarrhea. You may need to drink special liquids with salt and sugar, called Oral Rehydration Therapy. • Talk to your health care team if your diarrhea does not improve after 24 hours of taking diarrhea medication or if you have diarrhea more than 7 times in one day. <p>Ask your health care team for the Diarrhea pamphlet for more information.</p>	<p>Talk to your health care team if no improvement after 24 hours of taking diarrhea medication or if severe (more than 7 times in one day)</p>

Other rare, but serious side effects are possible with this treatment.

If you have **any** of the following, talk to your cancer health care team or get emergency medical help right away:

- Sudden changes to vision, speech, or the use of your arms or legs
- New pain in your chest, belly, arms, or legs
- Pain, swelling or hardening of a vein
- Irregular heartbeat, fainting, swelling in your legs, ankles or belly
- Sudden start of coughing, breathing problems, or coughing blood
- Dizziness or passing out, sudden, unexpected weight gain
- Severe headache, loss of consciousness, seizures, confusion
- Changes in personality, memory problems, not being able to speak properly
- New pain, numbness or tingling of your hands or feet, having trouble doing tasks (like doing up buttons, writing, walking) or if you have severe pain or numbness.
- Changes in your eyes (feeling painful, sensitive to light, blurry vision, seeing flashing lights or not able to see colours correctly)
- Severe rash, itchiness, swollen face, lip or tongue, chest or throat tightness during or shortly after the drug(s) are given
- Red or purple patches on the skin, rash/patches that do not turn white when pressed
- Red, itchy, bruised, and/or swollen skin, any burning, stinging or pain where the injection was given
- Red-brown coloured pee, severe tiredness or weakness, bruising easily

For more information on how to manage your symptoms ask your health care provider, or visit:

<https://www.cancercareontario.ca/symptoms>.

