

Regimen Monograph

[Regimen Name](#) | [Drug Regimen](#) | [Cycle Frequency](#) | [Premedication and Supportive Measures](#) | [Dose Modifications](#) | [Adverse Effects](#) | [Interactions](#) | [Drug Administration and Special Precautions](#) | [Recommended Clinical Monitoring](#) | [Administrative Information](#) | [References](#) | [Other Notes](#) | [Disclaimer](#)

A - Regimen Name

CRBPPEME Regimen

CARBOplatin-Pemetrexed**Disease Site** Lung - Non-Small Cell**Intent** Adjuvant**Regimen Category** **Evidence-Informed :**

Regimen is considered appropriate as part of the standard care of patients; meaningfully improves outcomes (survival, quality of life), tolerability or costs compared to alternatives (recommended by the Disease Site Team and national consensus body e.g. pan-Canadian Oncology Drug Review, pCODR). Recommendation is based on an appropriately conducted phase III clinical trial relevant to the Canadian context OR (where phase III trials are not feasible) an appropriately sized phase II trial. Regimens where one or more drugs are not approved by Health Canada for any indication will be identified under Rationale and Use.

Rationale and Uses For the treatment of completely resected stage II or IIIA non-small cell lung cancer with non-squamous histology.[back to top](#)

B - Drug Regimen

pemetrexed	500 mg /m ²	IV	Day 1
CARBOplatin	AUC 5*	IV	Day 1

*Adjust Carboplatin dose to AUC target (using Calvert formula) as outlined in "Other Notes" section.

[back to top](#)

C - Cycle Frequency**REPEAT EVERY 21 DAYS**

For 4 cycles unless disease progression or unacceptable toxicity occurs

[back to top](#)

D - Premedication and Supportive Measures

Antiemetic Regimen: Moderate + NK1 antagonist (Carboplatin AUC \geq 5)

Febrile Neutropenia Risk: Low

Other Supportive Care:

Also refer to [CCO Antiemetic Recommendations](#).

- Vitamin B12 1000mcg IM every 9 weeks, Folic acid 0.4 - 1 mg PO daily (both starting \geq 1 week prior to pemetrexed administration continue throughout and 3 weeks after last dose of Pemetrexed).
- Dexamethasone 4mg PO BID for 3 days starting day before chemotherapy suggested for rash prophylaxis.
- Note: NSAIDs should be held for 2-5 days prior and 2 days after pemetrexed (refer to pemetrexed monograph)

[back to top](#)

E - Dose Modifications

Doses should be modified according to the protocol by which the patient is being treated. The

following recommendations have been adapted from clinical trials or product monographs and could be considered.

Dosage with toxicity

Do not start a new cycle until ANC $\geq 1.5 \times 10^9/L$ and platelets $\geq 100 \times 10^9/L$. In the clinical trial, treatment delays of up to 3 weeks were permitted. Reduced doses were not re-escalated. Administration of G-CSF was permitted for febrile neutropenia, but was not a substitute for appropriate dose reductions (see table below).

Worse toxicity in previous cycle	Carboplatin (% previous dose)*	Pemetrexed (% previous dose)*
1 st episode febrile neutropenia, grade 4 thrombocytopenia or thrombocytopenic bleeding	75%	75%
2 nd episode febrile neutropenia, grade 4 thrombocytopenia or thrombocytopenic bleeding	50%**	50%**
Grade 3 related organ non-hematologic toxicity (except nausea and/or vomiting)	75%	75%
Grade 4 related organ non-hematologic toxicity	50% or Discontinue	50% or Discontinue
Symptoms suggesting pneumonitis	n/a	Hold and investigate; discontinue if confirmed.
Any occurrence of SJS/TEN	n/a	Discontinue

*Do not retreat unless platelets $\geq 100 \times 10^9/L$, ANC $\geq 1.5 \times 10^9/L$ and toxicities have recovered to \leq grade 1

**If toxicity recurs a 3rd time, discontinue

Hepatic Impairment

Pemetrexed is not extensively metabolized in the liver. No dosage adjustment is recommended, but use with caution. No dosage adjustment is recommended for carboplatin.

Renal Impairment

Patients with CrCl < 45 ml/min were excluded from clinical trials.

CrCl (ml/min)	Carboplatin dose	Pemetrexed dose
≥ 45 ml/min	Use Calvert formula, if appropriate	No change
< 45 ml/min	Discontinue	Discontinue

Dosage in the Elderly

Carboplatin: caution should be exercised and dose reduction considered as elderly patients may have more severe myelosuppression and neuropathy.

Pemetrexed: no dosage adjustments are needed, but elderly patients should be monitored closely as more myelosuppression, renal and severe GI effects were noted.

[back to top](#)

F - Adverse Effects

Refer to [pemetrexed](#), [CARBOplatin](#) drug monograph(s) for additional details of adverse effects.

Very common (≥ 50%)	Common (25-49%)	Less common (10-24%)	Uncommon (< 10%), but may be severe or life-threatening
<ul style="list-style-type: none"> • Myelosuppression ± infection, bleeding (may be severe) • Nausea, vomiting 	<ul style="list-style-type: none"> • Abnormal electrolyte(s) (↓ in Na, K, Ca, Mg) • Fatigue 	<ul style="list-style-type: none"> • Anorexia • Hearing impairment • Mucositis • Rash • Diarrhea • Nephrotoxicity (may be severe) • ↑ BUN • ↑ LFTs (transient) 	<ul style="list-style-type: none"> • Arrhythmia • Arterial / venous thromboembolism • Radiation recall reaction • GI perforation • Hemolytic uremic syndrome • Hemolytic anemia • Hemolysis • Peripheral ischemia • Hypersensitivity • Secondary

			malignancy <ul style="list-style-type: none"> • Peripheral neuropathy • Visual disorders • Encephalopathy • Pneumonitis
--	--	--	---

[back to top](#)

G - Interactions

Refer to [pemetrexed](#), [CARBOplatin](#) drug monograph(s) for additional details.

- Nephrotoxic drugs (e.g. aminoglycosides) may increase the toxicity of pemetrexed and exacerbate nephro and ototoxicity; caution and monitor closely if used together
- NSAIDs may increase the toxicity of pemetrexed. Hold NSAIDs with shorter half-lives (e.g. ibuprofen) at least 2 days before to 2 days after pemetrexed. Hold NSAIDs with long half-lives (e.g. piroxicam) 5 days before to 2 days after pemetrexed.
- Phenytoin levels may be altered by carboplatin. Monitor levels and adjust the phenytoin dose as required.
- Use with warfarin may increase INR and the risk of bleeding. Monitor INR and adjust the warfarin dose as required.

[back to top](#)

H - Drug Administration and Special Precautions

Refer to [pemetrexed](#), [CARBOplatin](#) drug monograph(s) for additional details.

Administration

Carboplatin:

- Mix in 100mL to 250mL bag (5% Dextrose or Normal Saline).
- Administer over 30 minutes, starting 30 minutes after the end of Pemetrexed.
- Incompatible with sets, needles or syringes containing aluminum – leads to precipitation and

loss of potency.

- Protect from light.

Pemetrexed:

- Reconstitute as directed with Normal Saline.
- Dilute drug in 100mL (Normal Saline only); Infuse IV over 10 minutes.
- Incompatible with calcium-containing solutions.
- Do not co-administer with other drugs and diluents.
- Keep unopened vials at room temperature. Pemetrexed is not light sensitive.

Contraindications

- Patients who have a hypersensitivity to these drugs or other platinum-containing compounds
- Patients with severe renal impairment (CrCl < 45 ml/min), severe myelosuppression or bleeding tumours
- Avoid the use of live vaccines

Warnings/precautions

- Patients with abnormal renal function or who are receiving concomitant nephrotoxic drugs
- Patients who have received extensive prior treatment, have poor performance status and those over 65 years of age
- Patients with cardiovascular risk factors

Pregnancy/lactation

- These drugs are not recommended in pregnancy.
- Adequate contraception should be used by both sexes during treatment, and for at least **6 months** after the last dose.
- Breastfeeding is not recommended

- Fertility Effects:
 - Carboplatin: Unknown
 - Pemetrexed: Yes (may be irreversible)

[back to top](#)

I - Recommended Clinical Monitoring

Treating physicians may decide to monitor more or less frequently for individual patients but should always consider recommendations from the product monograph.

Recommended Clinical Monitoring

- CBC; baseline and before each cycle
- Liver function tests; baseline and regular
- Renal function tests; baseline and regular, including electrolytes
- Clinical toxicity assessment for neurotoxicity, ototoxicity, hypersensitivity, bleeding, infection, GI, and pulmonary effects; at each visit
- Grade toxicity using the current [NCI-CTCAE \(Common Terminology Criteria for Adverse Events\) version](#)

Suggested Clinical Monitoring

INR for patients receiving warfarin; baseline and regular

[back to top](#)

J - Administrative Information

Approximate Patient Visit	2 hours
Pharmacy Workload (average time per visit)	33.069 minutes
Nursing Workload (average time per visit)	49.167 minutes

[back to top](#)

K - References

Gauvain C, Crequit P, Rousseau-Bussac G, et al. Adjuvant chemotherapy of non-small cell lung cancer: Tolerance of combined cisplatin-pemetrexed therapy. *Rev Mal Respir* 2014;31(9):817-21.

Kreuter M, Vansteenkiste J, Fischer JR, et al. Three-Year Follow-Up of a Randomized Phase II Trial on Refinement of Early-Stage NSCLC Adjuvant Chemotherapy with Cisplatin and Pemetrexed versus Cisplatin and Vinorelbine (the TREAT Study). *J Thorac Oncol* 2016;11(1):85-93.

Zhang L, Ou W, Liu Q, et al. Pemetrexed plus carboplatin as adjuvant chemotherapy in patients with curative resected non-squamous non-small cell lung cancer. *Thorac Cancer* 2014;5(1):50-6.

PEBC Advice Documents or Guidelines

- [Adjuvant Systemic and Radiation Therapy for Stage I to IIIA Completely Resected Non-Small-Cell Lung Cancers: ASCO-CCO Clinical Practice Guideline Update](#)

June 2021 removed pemetrexed NDFP funding info

[back to top](#)

L - Other Notes

Calvert Formula

DOSE (mg) = target AUC X (GFR + 25)

- AUC = product of serum concentration (mg/mL) and time (min)
- GFR (glomerular filtration rate) expressed as measured Creatinine Clearance or estimated from Serum Creatinine (by Cockcroft and Gault method or Jelliffe method)

(Calvert AH, Newell DR, Gumbrell LA, et al, Carboplatin dosage: Prospective evaluation of a simple formula based on renal function. *J Clin Oncol*, 1989; 7: 1748-1756)

[back to top](#)

M - Disclaimer

Regimen Abstracts

A Regimen Abstract is an abbreviated version of a Regimen Monograph and contains only top level information on usage, dosing, schedule, cycle length and special notes (if available). It is intended for healthcare providers and is to be used for informational purposes only. It is not intended to constitute or be a substitute for medical advice, and all

uses of the Regimen Abstract are subject to clinical judgment. Such information is provided on an “as-is” basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information’s quality, accuracy, currency, completeness, or reliability, and Cancer Care Ontario disclaims all liability for the use of this information, and for any claims, actions, demands or suits that arise from such use.

Information in regimen abstracts is accurate to the extent of the ST-QBP regimen master listings, and has not undergone the full review process of a regimen monograph. Full regimen monographs will be published for each ST-QBP regimen as they are developed.

Regimen Monographs

Refer to the [New Drug Funding Program](#) or [Ontario Public Drug Programs](#) websites for the most up-to-date public funding information.

The information set out in the drug monographs, regimen monographs, appendices and symptom management information (for health professionals) contained in the Drug Formulary (the “Formulary”) is intended for healthcare providers and is to be used for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or adverse effects of a particular drug, nor should it be construed to indicate that use of a particular drug is safe, appropriate or effective for a given condition. The information in the Formulary is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.

The format and content of the drug monographs, regimen monographs, appendices and symptom management information contained in the Formulary will change as they are reviewed and revised on a periodic basis. The date of last revision will be visible on each page of the monograph and regimen. Since standards of usage are constantly evolving, it is advised that the Formulary not be used as the sole source of information. It is strongly recommended that original references or product monograph be consulted prior to using a chemotherapy regimen for the first time.

Some Formulary documents, such as the medication information sheets, regimen information sheets and symptom management information (for patients), are intended for patients. Patients should always consult with their healthcare provider if they have questions regarding any information set out in the Formulary documents.

While care has been taken in the preparation of the information contained in the Formulary, such information is provided on an “as-is” basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information’s quality, accuracy, currency, completeness, or reliability.

CCO and the Formulary’s content providers shall have no liability, whether direct, indirect, consequential, contingent, special, or incidental, related to or arising from the information in the Formulary or its use thereof, whether based on breach of contract or tort (including negligence), and even if advised of the possibility thereof. Anyone using the information in the Formulary does so at his or her own risk, and by using such information, agrees to indemnify CCO and its content providers from any and all liability, loss, damages, costs and expenses (including legal fees and expenses) arising from such person’s use of the information in the Formulary.

[back to top](#)