

## Regimen Monograph

[Regimen Name](#) | [Drug Regimen](#) | [Cycle Frequency](#) | [Premedication and Supportive Measures](#) | [Dose Modifications](#) | [Adverse Effects](#) | [Interactions](#) | [Drug Administration and Special Precautions](#) | [Recommended Clinical Monitoring](#) | [Administrative Information](#) | [References](#) | [Other Notes](#) | [Disclaimer](#)

## A - Regimen Name

**DENO Regimen**

Denosumab

**Disease Site** Hematologic - Multiple Myeloma

**Intent** Palliative

**Regimen Category** **Evidence-Informed :**

Regimen is considered appropriate as part of the standard care of patients; meaningfully improves outcomes (survival, quality of life), tolerability or costs compared to alternatives (recommended by the Disease Site Team and national consensus body e.g. pan-Canadian Oncology Drug Review, pCODR). Recommendation is based on an appropriately conducted phase III clinical trial relevant to the Canadian context OR (where phase III trials are not feasible) an appropriately sized phase II trial. Regimens where one or more drugs are not approved by Health Canada for any indication will be identified under Rationale and Use.

**Rationale and Uses** For prevention of skeletal-related events in patients with newly diagnosed multiple myeloma.

[back to top](#)

## B - Drug Regimen

[denosumab](#) 120 mg Subcut Every 4 weeks

(This drug is not currently publicly funded for this regimen and intent)

[back to top](#)

---

**C - Cycle Frequency****REPEAT EVERY 28 DAYS**

Unless unacceptable toxicity

[back to top](#)

**D - Premedication and Supportive Measures**

**Antiemetic Regimen:** Not applicable

**Other Supportive Care:**

All patients, except those with hypercalcemia, should receive the following supplementation:

- at least 500mg of calcium daily
- at least 400 IU of vitamin D daily

[back to top](#)

**E - Dose Modifications**

Doses should be modified according to the protocol by which the patient is being treated. The following recommendations have been adapted from clinical trials or product monographs and could be considered.

Pre-existing hypocalcemia must be corrected prior to starting treatment.

Patients being treated with denosumab should not be treated concomitantly with bisphosphonates.

**Dosage with toxicity**

<b>Toxicity</b>	<b>Action</b>
Grade 3 or 4 drug-related toxicity	Consider holding or discontinuing
Osteonecrosis of the jaw	Follow guidelines for management. Consider holding or discontinuing treatment. Refer patient to dentist or oral surgeon.
Hypocalcemia	Treat appropriately. Consider holding or discontinuing treatment if severe.
Anaphylaxis or significant hypersensitivity	Treat appropriately. Discontinue denosumab permanently.

**Hepatic Impairment**

No studies have been conducted in patients with hepatic impairment.

**Renal Impairment**

No dose adjustment is required with renal impairment. Patients with renal impairment are at increased risk of severe life threatening hypocalcemia and require increased monitoring (refer to monitoring section).

**Dosage in the Elderly**

No adjustment required. No overall differences in safety and efficacy.

[back to top](#)

## F - Adverse Effects

Refer to [denosumab](#) drug monograph(s) for additional details of adverse effects

Common (25-49%)	Less common (10-24%)	Uncommon (< 10%), but may be severe or life-threatening
<ul style="list-style-type: none"> <li>• Nausea and vomiting</li> <li>• Fatigue</li> <li>• Anemia</li> <li>• Musculoskeletal pain</li> </ul>	<ul style="list-style-type: none"> <li>• Anorexia, weight loss</li> <li>• Constipation</li> <li>• Diarrhea</li> <li>• Abdominal pain</li> <li>• Cough, dyspnea (may be severe)</li> <li>• ↓ PO<sub>4</sub>, ↓ Ca (may be severe)</li> <li>• Headache</li> <li>• Fever</li> </ul>	<ul style="list-style-type: none"> <li>• Arterial thromboembolism</li> <li>• Venous thromboembolism</li> <li>• Renal failure</li> <li>• Arrhythmia, cardiotoxicity</li> <li>• Bone fracture (including atypical femoral; multiple vertebral after treatment discontinuation)</li> <li>• Osteonecrosis of the jaw</li> <li>• ↑ LFTs (may be severe)</li> <li>• Secondary malignancies</li> <li>• Hypersensitivity</li> </ul>

[back to top](#)

## G - Interactions

Refer to [denosumab](#) drug monograph(s) for additional details

- Exercise caution when given with drugs that may cause hypocalcemia. Monitor calcium levels closely.

[back to top](#)

## H - Drug Administration and Special Precautions

Refer to [denosumab](#) drug monograph(s) for additional details

**Administration:**

- Inject subcutaneously in the upper arm, upper thigh, or the abdomen.
- Should not be administered intravenously, intramuscularly or intradermally
- Use a 27-gauge needle to withdraw or inject the drug. Avoid vigorous shaking of the drug.
- Denosumab should appear clear, colourless to slightly yellow. It may contain trace amounts of translucent or white proteinaceous particles. Do not use if the solution is discoloured, cloudy, contains many particles or foreign matter.
- If a dose is missed, it may be given as soon as possible and the subsequent injection should be scheduled q4 weeks from the most recent injection date.
- Keep refrigerated in the original carton between 2-8°C. Protect from direct light
- Before use, the drug vial (in its original container) can be brought to room temperature (usually takes 15-30 minutes). Do not warm the drug by other methods. Once removed from the refrigerator, it must be stored at room temperature ( $\leq 25^{\circ}\text{C}$ ) and used within 30 days.

**Contraindications:**

- Contraindicated in patients who have a hypersensitivity to this drug or any of its components
- Do not use Xgeva® with Prolia®, as both products contain the same active ingredient, denosumab.

**Precautions:**

- Patients being treated with denosumab should not be treated concomitantly with bisphosphonates
- Pre-existing hypocalcemia must be corrected before starting denosumab treatment. Risk of hypocalcemia is greater in patients with moderate to severe renal impairment. Patients, except those with hypercalcemia, should receive adequate calcium and vitamin D supplementation (see Dosing section).
- Risk-benefit should be assessed for patients with risk factors for ONJ before starting treatment.
- Adequate contraception should be used by both sexes during treatment, at for at least 5 months after the last denosumab dose.
- Fertility effects: Unlikely

[back to top](#)

**I - Recommended Clinical Monitoring**

Treating physicians may decide to monitor more or less frequently for individual patients but should always consider recommendations from the product monograph.

---

### Recommended Clinical Monitoring

- Calcium, phosphate, magnesium - No hypercalcemia: baseline, within 2 weeks of the first dose, and as clinically indicated. In hypercalcemia: baseline, before each dose and as clinically indicated. Additional monitoring with renal dysfunction, symptoms of hypercalcemia and in patients with growing skeleton.
- Dental examination with appropriate preventative dentistry should be considered prior to treatment. Regular dental check-ups. Avoid invasive dental surgeries while on treatment.
- Vertebral fractures; evaluate patient risk after treatment discontinuation
- Clinical toxicity assessment for fatigue, musculoskeletal effects, hypocalcemia, ONJ, cough/dyspnea; at each visit
- Grade toxicity using the current [NCI-CTCAE \(Common Terminology Criteria for Adverse Events\) version](#)

[back to top](#)

### **J - Administrative Information**

Refer to local administration guidelines.

Pharmacy Workload (average time per visit) 13.85 minutes

Nursing Workload (average time per visit) 25.833 minutes

[back to top](#)

### **K - References**

Denosumab drug monograph, Cancer Care Ontario.

Raje N, Terpos E, Willenbacher W, et al. Denosumab versus zoledronic acid in bone disease treatment of newly diagnosed multiple myeloma: an international, double-blind, double-dummy, randomised, controlled, phase 3 study. *Lancet Oncol* 2018;19:370-81.

**October 2018** New ST-QBP regimen

[back to top](#)

## M - Disclaimer

### **Regimen Abstracts**

*A Regimen Abstract is an abbreviated version of a Regimen Monograph and contains only top level information on usage, dosing, schedule, cycle length and special notes (if available). It is intended for healthcare providers and is to be used for informational purposes only. It is not intended to constitute or be a substitute for medical advice, and all uses of the Regimen Abstract are subject to clinical judgment. Such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability, and Cancer Care Ontario disclaims all liability for the use of this information, and for any claims, actions, demands or suits that arise from such use.*

*Information in regimen abstracts is accurate to the extent of the ST-QBP regimen master listings, and has not undergone the full review process of a regimen monograph. Full regimen monographs will be published for each ST-QBP regimen as they are developed.*

### **Regimen Monographs**

Refer to the [New Drug Funding Program](#) or [Ontario Public Drug Programs](#) websites for the most up-to-date public funding information.

*The information set out in the drug monographs, regimen monographs, appendices and symptom management information (for health professionals) contained in the Drug Formulary (the "Formulary") is intended for healthcare providers and is to be used for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or adverse effects of a particular drug, nor should it be construed to indicate that use of a particular drug is safe, appropriate or effective for a given condition. The information in the Formulary is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.*

*The format and content of the drug monographs, regimen monographs, appendices and symptom management information contained in the Formulary will change as they are reviewed and revised on a periodic basis. The date of last revision will be visible on each page of the monograph and regimen. Since standards of usage are constantly evolving, it is advised that the Formulary not be used as the sole source of information. It is strongly recommended that original references or product monograph be consulted prior to using a chemotherapy regimen for the first time.*

*Some Formulary documents, such as the medication information sheets, regimen information sheets and symptom management information (for patients), are intended for patients. Patients should always consult with their healthcare provider if they have questions regarding any information set out in the Formulary documents.*

*While care has been taken in the preparation of the information contained in the Formulary, such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability.*

*CCO and the Formulary's content providers shall have no liability, whether direct, indirect, consequential, contingent, special, or incidental, related to or arising from the information in the Formulary or its use thereof, whether based on breach of contract or tort (including negligence), and even if advised of the possibility thereof. Anyone using the information in the Formulary does so at his or her own risk, and by using such information, agrees to indemnify CCO and its content providers from any and all liability, loss, damages, costs and expenses (including legal fees and expenses) arising from such person's use of the information in the Formulary.*

[back to top](#)