

## Regimen Monograph

[Regimen Name](#) | [Drug Regimen](#) | [Cycle Frequency](#) | [Premedication and Supportive Measures](#) | [Administrative Information](#) | [References](#) | [Other Notes](#) | [Disclaimer](#)

## A - Regimen Name

## FLUD+OBIN Regimen

Fludarabine-oBINutuzumab

**Disease Site** Hematologic - Lymphoma - Non-Hodgkin's Low Grade

**Intent** Palliative

**Regimen Category** **Evidence-informed :**

Regimen is considered appropriate as part of the standard care of patients; meaningfully improves outcomes (survival, quality of life), tolerability or costs compared to alternatives (recommended by the Disease Site Team and national consensus body e.g. pan-Canadian Oncology Drug Review, pCODR). Recommendation is based on an appropriately conducted phase III clinical trial relevant to the Canadian context OR (where phase III trials are not feasible) an appropriately sized phase II trial. Regimens where one or more drugs are not approved by Health Canada for any indication will be identified under Rationale and Use.

This **Regimen Abstract** is an **abbreviated** version of a Regimen Monograph and contains only top level information on usage, dosing, schedule, cycle length and special notes (if available). Information in regimen abstracts is accurate to the extent of the ST-QBP regimen master listings, and has not undergone the full review process of a regimen monograph. Full regimen monographs will be published for each ST-QBP regimen as they are developed.

**Rationale and Uses** For the treatment of patients with follicular lymphoma<sup>†</sup> whose disease is refractory\* to a rituximab-containing regimen and has a good performance status

<sup>†</sup> indolent lymphoma histologies other than follicular lymphoma (excluding CLL and mantle cell lymphoma) may be eligible for obinutuzumab funding (refer to [NDFP form](#))

\* no response to OR progression during or within 6 months after rituximab or a

---

 rituximab-containing regimen

**Supplementary Public Funding** [oBINutuzumab](#)  
 New Drug Funding Program (Obinutuzumab - In Combination with Chemotherapy for Refractory Follicular Lymphoma)

[back to top](#)

## B - Drug Regimen

### FLUD+OBIN (induction)

#### Cycle 1:

<a href="#">oBINutuzumab</a>	1000 mg	IV	Days 1, 8 and 15
<a href="#">fludarabine</a>	25 mg /m <sup>2</sup>	IV	Days 1 to 5

#### Cycles 2 to 6:

<a href="#">oBINutuzumab</a>	1000 mg	IV	Day 1
<a href="#">fludarabine</a>	25 mg /m <sup>2</sup>	IV	Days 1 to 5

For obinutuzumab maintenance use, report as regimen OBIN(MNT) after FLUD+OBIN induction.

[back to top](#)

## C - Cycle Frequency

**Induction: REPEAT EVERY 28 DAYS** for up to 6 cycles unless disease progression or unacceptable toxicity (see [NDFP form](#))

For patients who responded to induction therapy, refer to maintenance obinutuzumab regimen - OBIN(MNT).

[back to top](#)

**D - Premedication and Supportive Measures**

**Antiemetic Regimen:** Minimal

**Other Supportive Care:**

Also refer to [CCO Antiemetic Recommendations](#).

**Obinutuzumab:**

Hepatitis B screening should be performed prior to treatment for all patients.

Patients at risk for tumour lysis syndrome should receive adequate hydration and uricostatics or alternative starting 12 to 24 hours prior to infusion.

Consider withholding antihypertensives (if applicable) 12 hours prior to infusion, during infusion and for the first hour after drug administration, and withholding concomitant medications that increase bleeding risk, especially in the first cycle.

Patients with neutropenia should receive antimicrobial prophylaxis; consider use of G-CSF, antiviral and antifungal prophylaxis.

**Premedication recommendations:**

<b>Treatment cycle, day</b>	<b>Patients</b>	<b>Premedication</b>
Cycle 1, Day 1	All	IV corticosteroid <sup>*/^</sup> completed at least 1 hr prior to infusion &  PO analgesic/antipyretic <sup>**</sup> & antihistamine <sup>***</sup> at least 30 min prior to infusion
Subsequent infusions	Patients with no prior IR during previous infusion	PO analgesic/antipyretic <sup>**</sup> at least 30 min prior to infusion
	Patients with grade 1 or 2 IR with previous infusion	PO analgesic/antipyretic <sup>**</sup> & antihistamine <sup>***</sup> at least 30 min prior to infusion

	<p>Patients with grade 3 IR with previous infusion OR</p> <p>patients with lymphocyte counts &gt; 25 x 10<sup>9</sup>/L prior to next treatment</p>	<p>IV corticosteroid*/^ completed at least 1 hr prior to infusion &amp;</p> <p>PO analgesic/antipyretic** &amp; antihistamine*** at least 30 min prior to infusion</p>
--	---	--

\*e.g. 100 mg prednisone or 20 mg dexamethasone. Hydrocortisone should not be used as it has not been effective in reducing IR rates.

^ If a corticosteroid-containing chemotherapy regimen is given on the same day as obinutuzumab, the corticosteroid can be given as PO if given at least 1 hour prior to obinutuzumab, in which case additional IV corticosteroid as premedication is not required.

\*\*e.g. 1000 mg acetaminophen

\*\*\*e.g. 50 mg diphenhydramine

[back to top](#)

**J - Administrative Information**

Approximate Patient Visit	3.5 hours
Pharmacy Workload (average time per visit)	9.608 minutes
Nursing Workload (average time per visit)	43.967 minutes

[back to top](#)

## K - References

Cheson BD, Chua N, Mayer J, et al. Overall survival benefit in patients with rituximab-refractory indolent non-Hodgkin lymphoma who received obinutuzumab plus bendamustine induction and obinutuzumab maintenance in the GADOLIN study. *J Clin Oncol*. 2018 Aug 1;36(22):2259-66.

Coiffier B, Neidhardt-Berard EM, Tilly H, et al. Fludarabine alone compared to CHVP plus interferon in elderly patients with follicular lymphoma and adverse prognostic parameters: a GELA study. *Annals of Oncology* 1999; 10: 1191-7.

Klasa R, Meyer R, Shustik C, et al. Randomized phase III study of fludarabine phosphate versus cyclophosphamide, vincristine, and prednisone in patients with recurrent low-grade non-Hodgkin's lymphoma previously treated with an alkylating agent or alkylator-containing regimen. *J Clin Oncol* 2002; 20(24): 4649-54.

Radford J, Davies A, Cartron G, et al. Obinutuzumab (GA101) plus CHOP or FC in relapsed/refractory follicular lymphoma: results of the GAUDI study (BO21000). *Blood*. 2013 Aug 15;122(7):1137-43.

**June 2021** removed fludarabine NDFP funding info

[back to top](#)

## M - Disclaimer

### **Regimen Abstracts**

*A Regimen Abstract is an abbreviated version of a Regimen Monograph and contains only top level information on usage, dosing, schedule, cycle length and special notes (if available). It is intended for healthcare providers and is to be used for informational purposes only. It is not intended to constitute or be a substitute for medical advice, and all uses of the Regimen Abstract are subject to clinical judgment. Such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability, and Cancer Care Ontario disclaims all liability for the use of this information, and for any claims, actions, demands or suits that arise from such use.*

*Information in regimen abstracts is accurate to the extent of the ST-QBP regimen master listings, and has not undergone the full review process of a regimen monograph. Full regimen monographs will be published for each ST-QBP regimen as they are developed.*

### **Regimen Monographs**

*Refer to the [New Drug Funding Program](#) or [Ontario Public Drug Programs](#) websites for the most up-to-date public funding information.*

*The information set out in the drug monographs, regimen monographs, appendices and symptom management information (for health professionals) contained in the Drug Formulary (the "Formulary") is intended for healthcare*

*providers and is to be used for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or adverse effects of a particular drug, nor should it be construed to indicate that use of a particular drug is safe, appropriate or effective for a given condition. The information in the Formulary is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.*

*The format and content of the drug monographs, regimen monographs, appendices and symptom management information contained in the Formulary will change as they are reviewed and revised on a periodic basis. The date of last revision will be visible on each page of the monograph and regimen. Since standards of usage are constantly evolving, it is advised that the Formulary not be used as the sole source of information. It is strongly recommended that original references or product monograph be consulted prior to using a chemotherapy regimen for the first time.*

*Some Formulary documents, such as the medication information sheets, regimen information sheets and symptom management information (for patients), are intended for patients. Patients should always consult with their healthcare provider if they have questions regarding any information set out in the Formulary documents.*

*While care has been taken in the preparation of the information contained in the Formulary, such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability.*

*CCO and the Formulary's content providers shall have no liability, whether direct, indirect, consequential, contingent, special, or incidental, related to or arising from the information in the Formulary or its use thereof, whether based on breach of contract or tort (including negligence), and even if advised of the possibility thereof. Anyone using the information in the Formulary does so at his or her own risk, and by using such information, agrees to indemnify CCO and its content providers from any and all liability, loss, damages, costs and expenses (including legal fees and expenses) arising from such person's use of the information in the Formulary.*

[back to top](#)