

Regimen Monograph

[Regimen Name](#) | [Drug Regimen](#) | [Cycle Frequency](#) | [Premedication and Supportive Measures](#) | [Dose Modifications](#) | [Adverse Effects](#) | [Interactions](#) | [Drug Administration and Special Precautions](#) | [Recommended Clinical Monitoring](#) | [Administrative Information](#) | [References](#) | [Other Notes](#) | [Disclaimer](#)

A - Regimen Name

GEMC Regimen

Gemcitabine

Disease Site Lung - Mesothelioma

Intent Palliative

Regimen Category **Evidence-Informed :**

Regimen is considered appropriate as part of the standard care of patients; meaningfully improves outcomes (survival, quality of life), tolerability or costs compared to alternatives (recommended by the Disease Site Team and national consensus body e.g. pan-Canadian Oncology Drug Review, pCODR). Recommendation is based on an appropriately conducted phase III clinical trial relevant to the Canadian context OR (where phase III trials are not feasible) an appropriately sized phase II trial. Regimens where one or more drugs are not approved by Health Canada for any indication will be identified under Rationale and Use.

Rationale and Uses An alternative to pemetrexed-based therapy. GEMC should not be used in the second-line setting.

[back to top](#)

B - Drug Regimen

[gemcitabine](#) 1250 mg /m² IV Days 1, 8, 15

[back to top](#)

C - Cycle Frequency**REPEAT EVERY 28 DAYS**

For up to 10 cycles unless disease progression or unacceptable toxicity occurs

[back to top](#)

D - Premedication and Supportive Measures

Antiemetic Regimen: Low

Also refer to [CCO Antiemetic Recommendations](#).

[back to top](#)

E - Dose Modifications

Doses should be modified according to the protocol by which the patient is being treated.

Dosage with toxicity

Doses should not be re-escalated if they are reduced for non-hematological toxicities, febrile neutropenia or thrombocytopenic bleeding.

Table 1 - Day 1 of Cycle:

Worst Toxicity in Previous Cycle	% Full Dose
Non-hematologic Grade 3**	75%*
Non-hematologic Grade 4	Consider discontinuing, or 50-75%*
Febrile neutropenia, thrombocytopenic bleeding	75%*
> 1 Occurrence of Day 8/15 holds	75%*
<ul style="list-style-type: none"> • Pneumonitis • Hemolytic Uremic Syndrome (HUS) • Stevens-Johnson syndrome (SJS) • Toxic epidermal necrolysis (TEN) • Capillary Leak Syndrome (CLS) 	Discontinue

- Posterior reversible encephalopathy syndrome (PRES)

* Do not start new cycle until ANC $\geq 1.5 \times 10^9/L$, platelets $\geq 100 \times 10^9/L$ and non-hematologic toxicity \leq grade 2. Discontinue if non-hematological toxicities require more than a 50% dose reduction from the starting dose.

** except nausea/vomiting or alopecia

Other treatment days within cycle:

Table 2 - Non-hematologic toxicities

Toxicity	Action (% Full dose)
Grade 3**	HOLD; restart at 50-75%*
Grade 4	Discontinue

* Treat only if non-hematologic toxicities recover to \leq grade 2 and hematologic parameters are met on treatment day (Table 3). Discontinue if non-hematological toxicities require more than a 50% dose reduction from the starting dose.

** except nausea/vomiting, alopecia

Table 3 - Hematologic Toxicities:

Platelets on treatment day (x $10^9/L$)		ANC on treatment day (x $10^9/L$)	Action (% Full Dose)
>100	And	> 1	100% *
50 to 100	And/or	0.5 to 1	75% or consider omit*
<50	And/or	<0.5	Omit

* Treat only if above parameters are met on treatment day and non-hematologic toxicities \leq grade 2.

Hepatic Impairment

Gemcitabine should be used with caution in patients with hepatic impairment (cirrhosis, hepatitis, alcoholism, metastases, etc.); initial dose reduction should be considered if the patient is treated, especially in hyperbilirubinemia.

Suggested:

Bilirubin (micromol/L)	Starting dose
> 1.2 x ULN	800 mg/m ² ; escalate if tolerated

Renal Impairment

Gemcitabine should be used with caution in patients with renal insufficiency. There is insufficient information from clinical studies to allow clear dose recommendations for this patient population. Clinical trials with cisplatin mandated CrCl \geq 60mL/min. For patients with pre-existing renal insufficiency, the close monitoring for occurrence of hemolytic uremic syndrome is required.

Dosage in the Elderly

Decreased clearance and increased half-life occurs with increasing age, however no dose adjustment is necessary.

Dosage based on Gender

Decreased volume of distribution and clearance are seen in women, however no dose adjustment is necessary.

[back to top](#)

F - Adverse Effects

Refer to [gemcitabine](#) drug monograph(s) for additional details of adverse effects

Very common (\geq 50%)	Common (25-49%)	Less common (10-24%)	Uncommon (< 10%), but may be severe or life-threatening
---------------------------	-----------------	----------------------	---

<ul style="list-style-type: none"> • Myelosuppression ± infection, bleeding (may be severe) • ↑ LFTs • Nausea/ vomiting (generally mild) 	<ul style="list-style-type: none"> • Flu-like symptoms • Proteinuria • Rash (rarely severe) 	<ul style="list-style-type: none"> • Edema • Musculoskeletal pain • Alopecia (generally mild) • Diarrhea 	<ul style="list-style-type: none"> • Arrhythmia • Arterial thromboembolism • Cardiotoxicity • Hepatotoxicity including liver failure • Hemolytic-uremic syndrome • Creatinine increased • Hypersensitivity • Injection site reaction • Gangrene • RPLS/PRES • ILD/ARDS • Capillary leak syndrome • Vasculitis • Radiosensitization • Toxic epidermal necrolysis (TEN) • Stevens Johnson syndrome (SJS))
---	--	--	---

[back to top](#)

G - Interactions

Refer to [gemcitabine](#) drug monograph(s) for additional details

- No specific drug interaction studies have been conducted.
- Monitor INR closely with concurrent warfarin use and adjust warfarin dose as needed, as gemcitabine may decrease metabolism and synthesis of clotting factors.
- Gemcitabine is a known radiosensitizer.

[back to top](#)

H - Drug Administration and Special Precautions

Refer to [gemcitabine](#) drug monograph(s) for additional details

Administration

- May dilute reconstituted drug in normal saline for IV infusion, resulting in a minimum final concentration of at least 0.1 mg/mL.
- Gemcitabine is for IV administration only and should be infused over 30 minutes.
- To prevent increased toxicity, avoid an infusion time of > 60 minutes or dosing more frequently than once weekly

Contraindications

- Patients who have a hypersensitivity to this drug or any of its components.

Other Warnings/Precautions

- Use with extreme caution in patients with compromised bone marrow reserve.
- Use with caution in patients with hepatic impairment (including concurrent liver metastases or a previous history of hepatitis, alcoholism or liver cirrhosis) and patients with renal impairment.
- Acute shortness of breath with a temporal relationship to gemcitabine injection administration may occur.
- Patients receiving concurrent radiation while receiving the full dose gemcitabine should be closely monitored for reactions. Exacerbation of radiation therapy toxicity including potentially life-threatening esophagitis and pneumonitis, particularly in patients receiving large volumes of radiotherapy have been observed.

Pregnancy/Lactation

- Gemcitabine is not recommended for use in pregnancy. Adequate contraception should be used by both sexes during treatment, and for at least 6 months (general recommendation) after the last dose.
- Breastfeeding is not recommended.
- Fertility: Observed in animal studies
 - ◊ Decreased spermatogenesis and fertility in male mice.

[back to top](#)

I - Recommended Clinical Monitoring

Treating physicians may decide to monitor more or less frequently for individual patients but should always consider recommendations from the product monograph.

Recommended Clinical Monitoring

- CBC; Baseline and before each dose
- Liver function tests; Baseline, before each cycle and as clinically indicated
- Renal function tests; Baseline, before each cycle and as clinically indicated
- Clinical assessment of bleeding, infection, rash, diarrhea, nausea/vomiting, edema, injection site reactions, flu-like symptoms, hemolysis, signs/symptoms of capillary leak syndrome, cardiovascular, CNS and respiratory effects; At each visit
- Grade toxicity using the current [NCI-CTCAE \(Common Terminology Criteria for Adverse Events\) version](#)

Suggested Clinical Monitoring

- Urinalysis; baseline and as clinically indicated
- INR for patient receiving warfarin; baseline and as clinically indicated

[back to top](#)

J - Administrative Information

Approximate Patient Visit	0.75 hour
Pharmacy Workload (average time per visit)	22.855 minutes
Nursing Workload (average time per visit)	36.667 minutes

[back to top](#)

K - References

Gemcitabine drug monograph, Cancer Care Ontario

van Meerbeeck JP, Baas P, Debruyne C, et al. A Phase II study of gemcitabine in patients with malignant pleural mesothelioma. European Organization for Research and treatment of Cancer Lung

Cancer Cooperative Group. Cancer 1999;85(12):2577-82.

May 2020 Updated Adverse effects, Dosage with Toxicity, Drug Administration and Special Precautions and Recommended Clinical Monitoring sections.

[back to top](#)

M - Disclaimer

Regimen Abstracts

A Regimen Abstract is an abbreviated version of a Regimen Monograph and contains only top level information on usage, dosing, schedule, cycle length and special notes (if available). It is intended for healthcare providers and is to be used for informational purposes only. It is not intended to constitute or be a substitute for medical advice, and all uses of the Regimen Abstract are subject to clinical judgment. Such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability, and Cancer Care Ontario disclaims all liability for the use of this information, and for any claims, actions, demands or suits that arise from such use.

Information in regimen abstracts is accurate to the extent of the ST-QBP regimen master listings, and has not undergone the full review process of a regimen monograph. Full regimen monographs will be published for each ST-QBP regimen as they are developed.

Regimen Monographs

Refer to the [New Drug Funding Program](#) or [Ontario Public Drug Programs](#) websites for the most up-to-date public funding information.

The information set out in the drug monographs, regimen monographs, appendices and symptom management information (for health professionals) contained in the Drug Formulary (the "Formulary") is intended for healthcare providers and is to be used for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or adverse effects of a particular drug, nor should it be construed to indicate that use of a particular drug is safe, appropriate or effective for a given condition. The information in the Formulary is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.

The format and content of the drug monographs, regimen monographs, appendices and symptom management information contained in the Formulary will change as they are reviewed and revised on a periodic basis. The date of last revision will be visible on each page of the monograph and regimen. Since standards of usage are constantly evolving, it is advised that the Formulary not be used as the sole source of information. It is strongly recommended that original references or product monograph be consulted prior to using a chemotherapy regimen for the first time.

Some Formulary documents, such as the medication information sheets, regimen information sheets and symptom management information (for patients), are intended for patients. Patients should always consult with their healthcare provider if they have questions regarding any information set out in the Formulary documents.

While care has been taken in the preparation of the information contained in the Formulary, such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability.

CCO and the Formulary's content providers shall have no liability, whether direct, indirect, consequential, contingent, special, or incidental, related to or arising from the information in the Formulary or its use thereof, whether based on breach of contract or tort (including negligence), and even if advised of the possibility thereof. Anyone using the information in the Formulary does so at his or her own risk, and by using such information, agrees to indemnify CCO and its content providers from any and all liability, loss, damages, costs and expenses (including legal fees and expenses) arising from such person's use of the information in the Formulary.

[back to top](#)