

Medication Information Sheet

Kadcyla® trastuzumab emtansine

(tras-TOO-zue-mab em-

TAN-seen)

This document provides general information about your medication. It does not replace the advice of your health care professional. Always discuss your therapy with your health care professional and refer to the package insert for more details.

Other Name: T-DM1

Appearance: clear, colourless solution

What is this medication for?

- For treating certain types of breast cancer

What should I do before I have this medication?

- Tell your health care team if you have or had significant medical condition(s), especially if you have / had:
 - ◊ heart, liver or lung problems
 - ◊ breathing problems when you are at rest
 - ◊ problems with bleeding
 - ◊ nerve problems, such as numbness and tingling in your hands and feet, or
 - ◊ any allergies.
- Tell your health care team if you ever had serious reactions when being treated with trastuzumab products (such as Herceptin®, Herzuma®, Trazimera®, Kanjinti®, Ogivri®).
- Your health care team may do tests to examine your heart function before starting treatment. This is done to make sure it is safe for you to receive Kadcyla® trastuzumab emtansine.

Remember to:

- Tell your health care team about all of the other medications you are taking.
- Keep taking other medications that have been prescribed for you, unless you have been told not to by your health care team.

How will this medication affect sex, pregnancy and breastfeeding?

Talk to your health care team about:

- How this medication may affect your sexual health.
- How this medication may affect your ability to have a baby, if this applies to you.

This medication may harm an unborn baby. Tell your health care team if you or your partner are pregnant, become pregnant during treatment, or are breastfeeding.

- If there is any chance of pregnancy happening, you and your partner together must use **2 effective forms of birth control** at the same time until **7 months** after your last treatment dose.
- Do not use hormonal birth control (such as birth control pills), unless your health care team told you that they are safe. Talk to your health care team about the safest birth control for you.
- Do not breastfeed while taking this medication and for **7 weeks** after your last dose.

How is this medication given?

- This drug is given through an IV (injected into a vein) .
- It is usually given every 21 days. Talk to your health care team about your treatment schedule.
- This drug will be given (infused) over a longer period of time for the first cycle. If this goes well, it will be given over a shorter time for the rest of the cycles.
- If you missed your treatment appointment, talk to your health care team to find out what to do.

What else do I need to know while on this medication?

- **Will this medication interact with other medications or natural health products?**
 - ◊ This medication can interact with other medications, vitamins, foods and natural health products. Interactions can make the treatment not work as well or cause severe side effects.
 - ◊ Tell your health care team about all of your:
 - prescription and over-the-counter (non-prescription) medications and all other drugs, such as marijuana (medical or recreational)
 - natural health products such as vitamins, herbal teas, homeopathic medicines, and other supplements
 - ◊ Check with your health care team before starting or stopping any of them.

- **What should I do if I feel unwell, have pain, a headache or a fever?**
 - **Always** check your temperature to see if you have a fever **before** taking any medications for fever or pain (such as acetaminophen (Tylenol®) or ibuprofen (Advil®)).
 - Fever can be a sign of infection that may need treatment right away.
 - If you take these medications before you check for fever, they may lower your temperature and you may not know you have an infection.

How to check for fever:

Keep a digital (electronic) thermometer at home and take your temperature if you feel hot or unwell (for example, chills, headache, mild pain).

- You have a fever if your temperature taken in your mouth (oral temperature) is:
 - 38.3°C (100.9°F) or higher at any time
- OR
- 38.0°C (100.4°F) or higher for at least one hour.

If you do have a fever:

- **Try to contact your health care team. If you are not able to talk to them for advice, you MUST get emergency medical help right away.**
- Ask your health care team for the [Fever](#) pamphlet for more information.

If you do not have a fever but have mild symptoms such as headache or mild pain:

- Ask your health care team about the right medication for you. **Acetaminophen (Tylenol®)** is a safe choice for most people.
- **Talk to your health care team before you start taking** Ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or ASA (Aspirin®), as they may increase your chance of bleeding or interact with your cancer treatment.
- Talk to your health care team if you already take **low dose aspirin** for a medical condition (such as a heart problem). It may still be safe to take.

What to DO while on this medication:

- **DO** check with your health care team before getting any vaccinations, surgery, dental work or other medical procedures.

- DO talk to your health care team about your risk of getting heart problems during and after completing this treatment.
- DO tell your health care team if you have any new pain, numbness or tingling of your hands or feet. This is especially important if you are having trouble doing tasks (like doing up buttons, writing, walking) or if you have severe pain or numbness.
- DO consider asking someone to drive you to and from the hospital on your treatment days. You may feel tired or dizzy after your treatment.

What NOT to DO while on this medication:

- DO NOT smoke or drink alcohol while on treatment without talking to your health care team first. Smoking and drinking can make side effects worse and make your treatment not work as well.

What are the side effects of this medication?

The following table lists side effects that you may have when getting trastuzumab emtansine. The table is set up to list the most common side effects first and the least common last. It is unlikely that you will have all of the side effects listed and you may have some that are not listed.

Read over the side effect table so that you know what to look for and when to get help. Refer to this table if you experience any side effects while on trastuzumab emtansine.

Very Common Side Effects (50 or more out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Fatigue</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Feeling of tiredness or low energy that lasts a long time and does not go away with rest or sleep. <p>What to do?</p> <ul style="list-style-type: none"> • Be active. Aim to get 30 minutes of moderate exercise (you are able to talk comfortably while exercising) on most days. • Check with your health care team before starting any new exercise. • Pace yourself, do not rush. Put off less important activities. Rest when you need to. • Ask family or friends to help you with things like housework, shopping, and child or pet care. 	<p>Talk to your health care team if it does not improve or if it is severe</p>

Very Common Side Effects (50 or more out of 100 people)	
Side effects and what to do	When to contact health care team
<ul style="list-style-type: none"> • Eat well and drink at least 6 to 8 glasses of water or other liquids every day (unless your health care team has told you to drink more or less). • Avoid driving or using machinery if you are feeling tired. <p>Ask your health care team for the Fatigue pamphlet for more information.</p>	

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Nausea and vomiting</p> <p>(Generally mild)</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Nausea is feeling like you need to throw up. You may also feel light-headed. • You may feel nausea within hours to days after your treatment. <p>What to do?</p> <p>To help prevent nausea:</p> <ul style="list-style-type: none"> • It is easier to prevent nausea than to treat it once it happens. • Drink clear liquids and have small meals. Get fresh air and rest. • Do not eat spicy, fried foods or foods with a strong smell. • Limit caffeine (like coffee, tea) and avoid alcohol. <p>If you have nausea or vomiting:</p> <ul style="list-style-type: none"> • Take your rescue (as-needed) anti-nausea medication(s) as prescribed. • Ask your health care team for the Nausea & Vomiting pamphlet for more information. • Talk to your health care team if: <ul style="list-style-type: none"> ◦ nausea lasts more than 48 hours ◦ vomiting lasts more than 24 hours or if it is severe 	<p>Talk to your healthcare team if nausea lasts more than 48 hours or vomiting lasts more than 24 hours or if it is severe</p>

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Liver problems</p> <p>Your health care team may check your liver function with a blood test. The liver changes do not usually cause any symptoms.</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Rarely, you may develop yellowish skin or eyes, unusually dark pee or pain on the right side of your belly. This may be severe. <p>What to do?</p> <p>If you have any symptoms of liver problems, get emergency medical help right away.</p>	<p>Get emergency medical help right away</p>
<p>Mild joint, muscle pain or cramps; Headache</p> <p>What to look for?</p> <ul style="list-style-type: none"> • New pain in your muscles or joints, muscle cramps, or feeling achy • Mild headache <p>What to do?</p> <ul style="list-style-type: none"> • Take pain medication (acetaminophen or opioids such as codeine, morphine, hydromorphone, oxycodone) as prescribed. • Read the above section: "What should I do if I feel unwell, have pain, a headache or a fever?" before taking acetaminophen (Tylenol®), ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or Aspirin. These medications may hide an infection that needs treatment or they may increase your risk of bleeding. • Rest often and try light exercise (such as walking) as it may help. <p>Ask your health care team for the Pain pamphlet for more information.</p>	<p>Talk to your health care team if it does not improve or if it is severe</p>
<p>Low neutrophils (white blood cells) in the blood (neutropenia)</p> <p>(May be severe)</p> <p>When neutrophils are low, you are at risk of getting an infection more easily. Ask your health care team for the Neutropenia (Low white blood cell count) pamphlet for more information.</p>	<p>If you have a fever, try to contact your health care team. If you are unable to talk to the team for advice, you MUST get emergency medical help right away.</p>

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>What to look for?</p> <ul style="list-style-type: none"> • If you feel hot or unwell (for example if you have chills or a new cough), you must check your temperature to see if you have a fever. • Do not take medications that treat a fever before you take your temperature (for example, Tylenol®, acetaminophen, Advil® or ibuprofen). • Do not eat or drink anything hot or cold right before taking your temperature. <p>You have a fever if your temperature taken in your mouth (oral temperature) is:</p> <ul style="list-style-type: none"> • 38.3°C (100.9°F) or higher at any time OR • 38.0°C (100.4°F) or higher for at least one hour. <p>What to do?</p> <p>If your health care team has told you that you have low neutrophils:</p> <ul style="list-style-type: none"> • Wash your hands often to prevent infection. • Check with your health care team before getting any vaccines, surgeries, medical procedures or visiting your dentist. • Keep a digital thermometer at home so you can easily check for a fever. <p>If you have a fever:</p> <p>If you have a fever, try to contact your health care team. If you are unable to talk to the team for advice, you must get emergency medical help right away.</p>	
<p>Low platelets in the blood</p> <p>(May be severe)</p> <p>When your platelets are low, you are at risk for bleeding and bruising. Ask your health care team for the Low Platelet Count pamphlet for more information.</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Watch for signs of bleeding: <ul style="list-style-type: none"> ◦ bleeding from your gums ◦ unusual or heavy nosebleeds ◦ bruising easily or more than normal ◦ black coloured stools (poo) or blood in your stools (poo) ◦ coughing up red or brown coloured mucus ◦ dizziness, constant headache or changes in your vision 	<p>Talk to your health care team if you have any signs of bleeding. If you have bleeding that doesn't stop or is severe (very heavy), you MUST get emergency help right away.</p>

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<ul style="list-style-type: none"> ◦ heavy vaginal bleeding ◦ red or pink coloured urine (pee) <p>What to do?</p> <p>If your health care team has told you that you have low platelets:</p> <ul style="list-style-type: none"> • Tell your pharmacist that your platelet count may be low before taking any prescriptions or over-the-counter medication. • Check with your healthcare team before you go to the dentist. • Take care of your mouth and use a soft toothbrush. • Try to prevent cuts and bruises. • Ask your health care team what activities are safe for you. • Your treatment may have to be delayed if you have low platelets. Your health care team may recommend a blood transfusion. <p>If you have signs of bleeding:</p> <ul style="list-style-type: none"> • If you have a small bleed, clean the area with soap and water or a saline (saltwater) rinse. Apply pressure for at least 10 minutes. <p>If you have bleeding that does not stop or is severe (very heavy), you must get emergency medical help right away.</p>	
<p>Neuropathy (Tingling, numb toes or fingers)</p> <p>(May be severe)</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Numbness or tingling of your fingers and toes may happen after starting your treatment. • Sometimes it can be painful and feel like burning sensation, which may be severe. <p>What to do?</p> <ul style="list-style-type: none"> • Talk to your health care team if you have symptoms of neuropathy. • Numbness and tingling may slowly get better after your treatment ends. <p>In rare cases, it may continue long after treatment ends. If you continue to have bothersome symptoms, talk to your health care team for advice.</p>	<p>Talk to your health care team, especially if you have trouble doing tasks like doing up buttons, writing, moving, or if you have severe pain or numbness</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Constipation</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Having bowel movements (going poo) less often than normal. • Small hard stools (poo) that look like pellets. • The need to push hard and strain to have any stool (poo) come out. • Stomach ache or cramps. • A bloated belly, feeling of fullness, or discomfort. • Leaking of watery stools (poo). • Lots of gas or burping. • Nausea or vomiting. <p>What to do?</p> <p>To help prevent constipation:</p> <ul style="list-style-type: none"> • Try to eat more fiber rich foods like fruits with skin, leafy greens and whole grains. • Drink at least 6 to 8 cups of liquids each day unless your health care team has told you to drink more or less. • Be Active. Exercise can help to keep you regular. • If you take opioid pain medication, ask your health care team if eating more fibre is right for you. <p>To help treat constipation:</p> <ul style="list-style-type: none"> • If you have not had a bowel movement in 2 to 3 days you may need to take a laxative (medication to help you poo) to help you have regular bowel movements. Ask your health care team what to do. <p>Ask your health care team for the Constipation Pamphlet for more information.</p>	<p>Talk to your health care team if it does not improve or if it is severe</p>
<p>Mouth sores</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Round, painful, white or gray sores inside your mouth that can occur on the tongue, lips, gums, or inside your cheeks. • In more severe cases they may make it hard to swallow, eat or brush your teeth. • They may last for 3 days or longer. 	<p>Talk to your health care team as soon as you notice mouth or lip sores or if it hurts to eat, drink or swallow</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>What to do?</p> <p>To help prevent mouth sores:</p> <ul style="list-style-type: none"> • Take care of your mouth by gently brushing and flossing regularly. • Rinse your mouth often with a homemade mouthwash. • To make a homemade mouthwash, mix 1 teaspoonful of baking soda and 1 teaspoonful of salt in 4 cups (1L) of water. • Do not use store-bought mouthwashes, especially those with alcohol, because they may irritate your mouth. <p>If you have mouth sores:</p> <ul style="list-style-type: none"> • Avoid hot, spicy, acidic, hard or crunchy foods. • Your doctor may prescribe a special mouthwash to relieve mouth sores and prevent infection. • Talk to your health care team as soon as you notice mouth or lip sores or if it hurts to eat, drink or swallow. <p>Ask your health care team for the Oral Care (Mouth Care) pamphlet for more information.</p>	
<p>Cough and feeling short of breath</p> <p>What to look for?</p> <ul style="list-style-type: none"> • You may have a cough and feel short of breath. • Symptoms that commonly occur with a cough are: <ul style="list-style-type: none"> ◦ Wheezing or a whistling breathing ◦ Runny nose ◦ Sore throat ◦ Heartburn ◦ Weight loss ◦ Fever and chills • Rarely this may be severe with chest pain, trouble breathing or coughing up blood. <p>What to do?</p> <ul style="list-style-type: none"> • Check your temperature to see if you have a fever. Read the above section "What should I do if I feel unwell, have pain, a headache or a fever?". • If you have a fever, try to talk to your health care team. If you are not able to talk to them for advice, you MUST get emergency medical help right away. 	<p>Talk to your health care team. If you are not able to talk to your health care team for advice, and you have a fever or severe symptoms, you MUST get emergency medical help right away</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<ul style="list-style-type: none"> If you have a severe cough with chest pain, trouble breathing or you are coughing up blood, get medical help right away. 	
<p>Dry mouth</p> <p>What to look for?</p> <ul style="list-style-type: none"> You may have a dry or sticky feeling in your mouth or throat, Your saliva may be thick and stringy. You may have cracks in your lips or at the corners of your mouth You may have difficulty chewing, tasting, swallowing or talking <p>What to do?</p> <ul style="list-style-type: none"> Use sugar-free gum or lozenges (e.g. those that contain xylitol) to help keep your mouth moist. Suck on ice chips or sugarless popsicles to help relieve dry mouth. Rinse your mouth often (every 1 to 2 hours) with a homemade mouthwash. To make a homemade mouthwash, mix 1 teaspoon of baking soda and 1 teaspoon of salt in 4 cups (1L) of water Do not use store-bought mouthwashes, especially those with alcohol because they may irritate dry your mouth Spray your mouth with water or artificial saliva products (e.g., Moi-Stir Spray®, Biotene® products) as needed to keep it moist. Apply mouth lubricant (like Biotene Oral balance gel®) after you brush your teeth, at bedtime, and as needed. Use a steam vaporizer at night to relieve nighttime dry mouth Speak to your health care team about the right product for you. <p>See our Mouth Care pamphlet for more information.</p> <p>Talk to your health care team if your dry mouth does not improve or if it is severe.</p>	<p>Talk to your health care team if it does not improve or if it is severe</p>
<p>Trouble Sleeping</p> <p>Your medications may cause trouble sleeping. It may get better once your body gets used to the medication or when your treatment ends.</p> <p>What to look for?</p> <ul style="list-style-type: none"> You may find it hard to fall asleep or stay asleep. How well you sleep may change over your treatment. For example, you may have 	<p>Talk to your health care team if it does not improve or if it is severe</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>several nights of poor sleep followed by a night of better sleep.</p> <ul style="list-style-type: none"> You may wake up too early or not feel well-rested after a night's sleep. You may feel tired or sleepy during the day. <p>What to do?</p> <p>Talk to your health care team if it does not improve or if it is severe</p>	
<p>Diarrhea</p> <p>What to look for?</p> <ul style="list-style-type: none"> Loose, watery, unformed stool (poo) that may happen days to weeks after you get your treatment. <p>What to do?</p> <p>If you have diarrhea:</p> <ul style="list-style-type: none"> Take anti-diarrhea medication if your health care team prescribed it or told you to take it. Do not eat foods or drinks with artificial sweetener (like chewing gum or 'diet' drinks), coffee and alcohol. Eat many small meals and snacks instead of 2 or 3 large meals. Drink at least 6 to 8 cups of liquids each day, unless your health care team has told you to drink more or less. Talk to your health care team if you can't drink 6 to 8 cups of liquids each day when you have diarrhea. You may need to drink special liquids with salt and sugar, called Oral Rehydration Therapy. Talk to your health care team if your diarrhea does not improve after 24 hours of taking diarrhea medication or if you have diarrhea more than 7 times in one day. <p>Ask your health care team for the Diarrhea pamphlet for more information.</p>	<p>Talk to your health care team if no improvement after 24 hours of taking diarrhea medication or if severe (more than 7 times in one day)</p>
<p>Dizziness</p> <p>What to look for?</p> <ul style="list-style-type: none"> You may feel light-headed and like you might faint (pass out). <p>What to do?</p> <ul style="list-style-type: none"> Lay down right away so you do not fall. 	<p>Talk to your health care team if it does not improve or if it is severe</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<ul style="list-style-type: none"> • Slowly get up and start moving once you feel better. • Do not drive a motor vehicle or use machinery if you feel dizzy. 	
<p>Reactions at the injection site</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Your skin may become red, itchy, bruised, and/or swollen where the injection was given. • Site reactions are usually mild and go away within one to three days. <p>What to do?</p> <ul style="list-style-type: none"> • Tell your nurse right away if there is any burning, stinging or other pain while trastuzumab emtansine is being injected into your vein. • Talk to your health care team if injection site reactions do not improve or are severe. 	<p>Talk to your health care team if it does not improve or if it is severe</p>

Other rare, but serious side effects are possible.

If you experience ANY of the following, speak to your cancer health care provider or get emergency medical help right away:

- Chest pain, fainting (passing out) or unexpected swelling in your legs, ankles and belly
- Symptoms of an allergic reaction (during or soon after the medication is given): flushing, fever, itchiness, rash, swollen lips, face or tongue, wheezing, throat and/or chest tightness
- Difficulty thinking or speaking, confusion, personality changes, forgetfulness, seizures
- Eye pain, sensitivity to light, blurred vision, or any changes to your vision

Who do I contact if I have questions or need help?

My cancer health care provider is: _____

During the day I should contact: _____

Evenings, weekends and holidays: _____

