

## Regimen Monograph

[Regimen Name](#) | [Drug Regimen](#) | [Cycle Frequency](#) | [Premedication and Supportive Measures](#) | [Dose Modifications](#) | [Adverse Effects](#) | [Interactions](#) | [Drug Administration and Special Precautions](#) | [Recommended Clinical Monitoring](#) | [Administrative Information](#) | [References](#) | [Other Notes](#) | [Disclaimer](#)

## A - Regimen Name

**ZOLE Regimen**

Zoledronic Acid

**Disease Site** Genitourinary  
Renal Cell / Kidney

**Intent** Palliative

**Regimen Category** **Evidence-Informed :**

Regimen is considered appropriate as part of the standard care of patients; meaningfully improves outcomes (survival, quality of life), tolerability or costs compared to alternatives (recommended by the Disease Site Team and national consensus body e.g. pan-Canadian Oncology Drug Review, pCODR). Recommendation is based on an appropriately conducted phase III clinical trial relevant to the Canadian context OR (where phase III trials are not feasible) an appropriately sized phase II trial. Regimens where one or more drugs are not approved by Health Canada for any indication will be identified under Rationale and Use.

**Rationale and Uses** For the treatment patients with renal cell cancer and bone metastases.

[back to top](#)

## B - Drug Regimen

[zoledronic acid](#) 4 mg IV Day 1

[back to top](#)

**C - Cycle Frequency****REPEAT EVERY 21 DAYS**

unless unacceptable toxicity.

[back to top](#)

**D - Premedication and Supportive Measures****Other Supportive Care:**

All patients, especially those with hypercalcemia, should be adequately hydrated. Calcium and Vitamin D supplements should be considered in patients who have normal calcium levels with no history of hypercalcemia. (Refer to zoledronic acid monograph).

[back to top](#)

**E - Dose Modifications**

Doses should be modified according to the protocol by which the patient is being treated. The following recommendations have been adapted from clinical trials or product monographs and could be considered.

Do not administer to patients with open soft tissue lesions in the mouth.

Hypocalcemia must be corrected before administering zoledronic acid.

**Dosage with toxicity**

Dosage in myelosuppression: No dosage adjustment required

<b>Toxicity</b>	<b>Action</b>
Atypical fractures of the femur	Hold if suspected. Consider discontinuing if confirmed.
Ocular symptoms other than uncomplicated conjunctivitis	Refer to ophthalmologist; consider discontinuing
Osteonecrosis of the jaw, other sites	For ONJ, refer to dentist or dental surgeon; consider hold or discontinue
Severe musculoskeletal pain	Discontinue
Acquired Fanconi syndrome	Discontinue
Increased creatinine:	Hold until recovered to within 10% of

<ol style="list-style-type: none"> <li>1. <math>\geq 44 \mu\text{mol/L}</math> <math>\uparrow</math> if normal baseline** OR</li> <li>2. <math>\geq 88 \mu\text{mol/L}</math> <math>\uparrow</math> if abnormal at baseline OR</li> <li>3. Serum creatinine <math>&gt; 265 \mu\text{mol/L}</math></li> </ol>	baseline (see table for dose adjustment for renal impairment at baseline)
--	---

\*\*normal baseline creatinine is defined as  $< 123 \mu\text{mol/L}$

### **Hepatic Impairment**

There are no pharmacokinetic data in patients with hepatic impairment. Zoledronic acid is not cleared by the liver.

### **Renal Impairment**

			Starting Dose
Creatinine		Creatinine Clearance (mL/min)	For bone metastases
		$> 60$	4 mg
		50 - 60	3.5 mg
		40 - 49	3.3 mg
		30 - 39	3 mg
$> 265 \mu\text{mol/L}$	Or	$< 30$	Do not treat

### **Dosage in the Elderly**

Similar efficacy and safety as compared to younger patients, but use with caution due to cardiac risks or renal function impairment.

[back to top](#)

**F - Adverse Effects**

Refer to [zoledronic acid](#) drug monograph(s) for additional details of adverse effects

<b>Common (25-49%)</b>	<b>Less common (10-24%)</b>	<b>Uncommon (&lt; 10%), but may be severe or life-threatening</b>
<ul style="list-style-type: none"> <li>• Nausea, vomiting</li> <li>• Fatigue, flu-like symptoms</li> <li>• Cough, dyspnea (may be severe)</li> </ul>	<ul style="list-style-type: none"> <li>• Diarrhea</li> <li>• Musculoskeletal pain (may be severe)</li> <li>• Edema</li> <li>• Headache</li> <li>• Dizziness</li> <li>• Nephrotoxicity (may be severe)</li> <li>• Weight loss</li> <li>• Paresthesia</li> <li>• Depression</li> <li>• Abnormal electrolytes</li> <li>• Conjunctivitis</li> </ul>	<ul style="list-style-type: none"> <li>• Atypical fractures of the femur</li> <li>• Atrial fibrillation, arrhythmia</li> <li>• Osteonecrosis of the jaw (ONJ) or other sites</li> <li>• Hypersensitivity</li> <li>• Eye disorders</li> <li>• Acquired Fanconi syndrome</li> </ul>

[back to top](#)

**G - Interactions**

Refer to [zoledronic acid](#) drug monograph(s) for additional details

- Caution and monitor with drugs that cause hypocalcemia (e.g. aminoglycosides, loop diuretics, calcitonin)
- Caution and monitor with drugs that cause renal dysfunction (e.g. NSAIDs, ACE inhibitors)
- Avoid in patients with hypersensitivity to ASA given possible increased risk of bronchospasm (theoretical)
- Caution with antiangiogenic drugs (e.g. sunitinib, bevacizumab) given increased risk of ONJ

[back to top](#)

---

## H - Drug Administration and Special Precautions

Refer to [zoledronic acid](#) drug monograph(s) for additional details

### Administration:

- Do not infuse over a duration of less than 15 minutes.
- All patients should be adequately hydrated prior to and after administration of zoledronic acid, but overhydration should be avoided.
- Mix with 100 mL solution (D5W or NS) and infuse over  $\geq 15$  minutes.
- Do not mix with calcium or other divalent cation-containing solutions.
- Compatible with PVC, glass, polyethylene and polypropylene containers or infusion lines.
- Should be administered as a single intravenous solution in a line separate from all other drugs.
- Store unopened vials at room temperature.

### Contraindications:

- Patients who have a hypersensitivity to this drug or any of its components, or other bisphosphonates
- Patients with non-corrected hypocalcemia at time of infusion or severe renal failure
- Zoledronic acid should not be given together with other bisphosphonates since the combined effects of these agents are unknown

### Other Warnings/Precautions:

- The use of zoledronic acid with other nephrotoxins, doses  $> 4$ mg, infusion duration  $< 15$  minutes and previous bisphosphonate use are associated with an increased risk of renal failure.
- Use with caution in patients with cardiac failure, especially in the elderly.
- Use with caution in patients with risk factors for ONJ, including patients receiving concomitant chemotherapy or anti-angiogenic agents; patients should be advised to avoid invasive dental procedures while receiving zoledronic acid.
- Caution in patients who have had thyroid surgery since they are susceptible to hypocalcaemia due to relative hypoparathyroidism.

[back to top](#)

## I - Recommended Clinical Monitoring

Treating physicians may decide to monitor more or less frequently for individual patients but should always consider recommendations from the product monograph.

### Recommended Clinical Monitoring

- Renal function tests (serum creatinine and BUN); baseline, before each dose and during therapy, as indicated
- Calcium, corrected levels (including serum albumin), electrolytes (including phosphate, magnesium); baseline, before each dose and during therapy, as indicated
- CBC; baseline and as clinically indicated
- Comprehensive dental evaluation of both hard and soft tissues before starting bisphosphonate treatment; undergo invasive dental procedures, if needed, before starting bisphosphonate treatment; regular check-ups
- Clinical toxicity assessment for flu-like syndrome, dental, signs of acquired Fanconi syndrome, musculoskeletal and ocular symptoms; at each visit
- Grade toxicity using the current [NCI-CTCAE \(Common Terminology Criteria for Adverse Events\) version](#)

### Suggested Clinical Monitoring

- Ophthalmology examination with ocular symptoms; as clinically indicated

[back to top](#)

## J - Administrative Information

Approximate Patient Visit	0.5 hour
Pharmacy Workload (average time per visit)	16 minutes
Nursing Workload (average time per visit)	35 minutes

[back to top](#)

## K - References

Lipton A, Zheng M, Seaman J. Zoledronic acid delays the onset of skeletal-related events and progression of skeletal disease in patients with advanced renal cell carcinoma. Cancer

---

2003;98(5):962-9.

Rosen LS, Gordon D, Tchekmedyian S, et al. Zoledronic acid versus placebo in the treatment of skeletal metastases in patients with lung cancer and other solid tumors: a phase III, double-blind, randomized trial--the zoledronic acid lung cancer and other solid tumors study group. *J Clin Oncol* 2003;21(16):3150-7.

Zoledronic acid drug monograph, Cancer Care Ontario.

**April 2023** removed NDFP form

[back to top](#)

## L - Other Notes

The skeletal-related event and skeletal morbidity rates were significantly less for patients who received zoledronic acid compared to placebo. Time to first skeletal-related event and time-to-progression of bone lesions were also significantly better in the zoledronic acid arm.

In a small retrospective review of 166 cancer patients, of whom 78 received zoledronic acid, osteonecrosis of the jaw (ONJ) developed in one patient with renal cell carcinoma who received sunitinib and zoledronic acid concurrently. A significantly higher incidence of ONJ was observed in patients with cancer receiving bisphosphonates with antiangiogenic agents versus bisphosphonates treatment alone (16% vs 1.1%, p=0.008).

[back to top](#)

## M - Disclaimer

### **Regimen Abstracts**

*A Regimen Abstract is an abbreviated version of a Regimen Monograph and contains only top level information on usage, dosing, schedule, cycle length and special notes (if available). It is intended for healthcare providers and is to be used for informational purposes only. It is not intended to constitute or be a substitute for medical advice, and all uses of the Regimen Abstract are subject to clinical judgment. Such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability, and Cancer Care Ontario disclaims all liability for the use of this information, and for any claims, actions, demands or suits that arise from such use.*

*Information in regimen abstracts is accurate to the extent of the ST-QBP regimen master listings, and has not undergone the full review process of a regimen monograph. Full regimen monographs will be published for each ST-QBP regimen as they are developed.*

---

**Regimen Monographs**

Refer to the [New Drug Funding Program](#) or [Ontario Public Drug Programs](#) websites for the most up-to-date public funding information.

*The information set out in the drug monographs, regimen monographs, appendices and symptom management information (for health professionals) contained in the Drug Formulary (the "Formulary") is intended for healthcare providers and is to be used for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or adverse effects of a particular drug, nor should it be construed to indicate that use of a particular drug is safe, appropriate or effective for a given condition. The information in the Formulary is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.*

*The format and content of the drug monographs, regimen monographs, appendices and symptom management information contained in the Formulary will change as they are reviewed and revised on a periodic basis. The date of last revision will be visible on each page of the monograph and regimen. Since standards of usage are constantly evolving, it is advised that the Formulary not be used as the sole source of information. It is strongly recommended that original references or product monograph be consulted prior to using a chemotherapy regimen for the first time.*

*Some Formulary documents, such as the medication information sheets, regimen information sheets and symptom management information (for patients), are intended for patients. Patients should always consult with their healthcare provider if they have questions regarding any information set out in the Formulary documents.*

*While care has been taken in the preparation of the information contained in the Formulary, such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability.*

*CCO and the Formulary's content providers shall have no liability, whether direct, indirect, consequential, contingent, special, or incidental, related to or arising from the information in the Formulary or its use thereof, whether based on breach of contract or tort (including negligence), and even if advised of the possibility thereof. Anyone using the information in the Formulary does so at his or her own risk, and by using such information, agrees to indemnify CCO and its content providers from any and all liability, loss, damages, costs and expenses (including legal fees and expenses) arising from such person's use of the information in the Formulary.*

[back to top](#)