

**Drug Monograph**

[Drug Name](#) | [Mechanism of Action and Pharmacokinetics](#) | [Indications and Status](#) | [Adverse Effects](#) | [Dosing](#) | [Administration Guidelines](#) | [Special Precautions](#) | [Interactions](#) | [Recommended Clinical Monitoring](#) | [Supplementary Public Funding](#) | [References](#) | [Disclaimer](#)

**A - Drug Name**

# aldesleukin

**SYNONYM(S):** interleukin-2; IL-2; lymphocyte mitogenic factor; T-cell growth factor; thymocyte stimulating factor

**COMMON TRADE NAME(S):** Proleukin® (interleukin-2)

[back to top](#)

**B - Mechanism of Action and Pharmacokinetics**

Aldesleukin, a recombinant interleukin-2, has multiple immunologic effects, including activation of cellular immunity with lymphocytosis, eosinophilia, thrombocytopenia and production of lymphokines including TNF, IL-1 and gamma-interferon. Tumour growth inhibition has been reported in *in vivo* studies. Objective responses are seen in  $\pm$  16% of patients but no overall survival benefit in controlled studies has been demonstrated.

Absorption	Oral: no	
Distribution	Pharmacokinetics appear to be dose proportional. Uptake into lung, liver, spleen and kidney is rapid.	
	Cross blood brain barrier?	No information found
	PPB	No information found
Metabolism	Aldesleukin is rapidly metabolized to its composite amino acids in the proximal tubules of the kidney.	
	Active metabolites	Trace
	Inactive metabolites	Yes (to its composite amino acids)

Elimination	2- compartmental disposition, cleared as inactive metabolites by glomerular filtration and peritubular extraction	
	Clearance	268mL/minute
	Half-life	13-85 minutes
	Urine	80% via renal tubular and glomerular filtration

[back to top](#)

## C - Indications and Status

### Health Canada Approvals:

- Carefully selected adults with metastatic renal cell carcinoma
- Carefully selected adults with metastatic malignant melanoma

[back to top](#)

## D - Adverse Effects

Minimal (Intralesional)  
Low (IV doses  $\leq$  12 MU/m<sup>2</sup>)

**Emetogenic Potential:** Moderate (IV doses >12-15 MU/m<sup>2</sup>)

**Extravasation Potential:** None

ORGAN SITE	SIDE EFFECT* (%)	ONSET**
Cardiovascular	Arterial thromboembolism (rare)	E
	Cardiotoxicity (<1%)	E
	Venous thromboembolism (rare)	E
	Ventricular arrhythmia (12%) (supraventricular)	I
Dermatological	Other (persistent vitiligo; rare)	E
	Rash (42%) (may be exfoliative)	I E
Gastrointestinal	Abdominal pain (11%)	E
	Anorexia (20%)	E

	Diarrhea (67%)	E
	GI perforation (or obstruction, +/- bleeding; rare)	E
	Mucositis (22%)	E
	Nausea, vomiting (50%)	I E
General	Fatigue (27%)	E
	Flu-like symptoms (52%)	I
	Hyperthermia (1%) (malignant hyper or hypothermia)	E
Hematological	Disseminated intravascular coagulation (1%)	I
	Myelosuppression ± infection, bleeding (37%) (severe 1%)	E
Hepatobiliary	↑ LFTs (40%) (may be severe)	E
	Pancreatitis (rare)	E
Hypersensitivity	Drug reaction (rare)	E
Immune	Other (risk of rejection in allograft recipients, exacerbation of autoimmune diseases)	E D
Infection	Infection (13%) (may be severe, including atypical)	E
Metabolic / Endocrine	Abnormal electrolyte(s) (12%) (decreased Ca, Mg)	E
	Acidosis (12%)	I E
Musculoskeletal	Rhabdomyolysis (especially with interferon; rare)	E
Nervous System	Confusion (34%) (includes somnolence)	E
	Depression (4%) (may be severe)	E D
	Dizziness (11%)	E
	Leukoencephalopathy (rare)	E
	Neuropathy (6%; optic neuritis, demyelination-rare)	E
	Seizure (<1%)	E
Ophthalmic	Conjunctivitis (2%)	E
	Eye disorders (1%)	E
Renal	Creatinine increased (33%) (may be severe)	E
	Tumor lysis syndrome (with chemotherapy)	E
Respiratory	Adult respiratory distress syndrome (ARDS) (3%)	E
	Dyspnea (43%)	E
Vascular	Capillary leak syndrome (71%)	I E
	Vasculitis (cutaneous; cerebral- rare)	E

---

\* "*Incidence*" may refer to an absolute value or the higher value from a reported range.  
"*Rare*" may refer to events with < 1% incidence, reported in post-marketing, phase 1 studies, isolated data or anecdotal reports.

\*\* I = *immediate* (onset in hours to days)    E = *early* (days to weeks)  
D = *delayed* (weeks to months)    L = *late* (months to years)

Most adverse reactions usually reverse or improve within 2-3 days of discontinuing therapy, due to aldesleukin's short half-life, but drug-related deaths occur in 2-4% of patients treated with aldesleukin.

The major dose-limiting side effect of aldesleukin is **capillary leak syndrome**. Aldesleukin administration induces decreased vascular tone and increased vascular permeability leading to hypotension, reduced organ perfusion and function (renal, cardiac, hepatic, etc.), ascites, effusions, cardiac arrhythmias/ischemia and respiratory insufficiency. The management of capillary leak syndrome involves careful monitoring of fluid and organ perfusion. Administration of intravenous dopamine or phenylephrine to increase blood pressure may help maintain organ perfusion, particularly renal perfusion and thereby preserve urine output. Administration of subsequent doses of aldesleukin should be delayed until recovery of organ perfusion is observed.

**Renal dysfunction** is reversible and also usually secondary to capillary leak syndrome. It is also correlated with the dose, duration of treatment and the patient's baseline renal function. The administration of indomethacin for flu-like syndrome may potentiate renal dysfunction by decreasing intrarenal prostaglandins.

A **flu-like syndrome (fever, rigors and chills)** develops in most patients receiving aldesleukin. Treatment with acetaminophen or a NSAID (e.g. ibuprofen, indomethacin) may minimize the risk, but renal function should be monitored carefully. In clinical trials, meperidine was administered to control the rigors associated with fever.

The clinical significance of developing **non-neutralizing anti-interleukin-2 antibodies** to aldesleukin is unknown. The incidence of antibodies formation is less than 1%. Exacerbation of autoimmune disorders such as Crohns, myasthenia, vasculitis, hypo or hyperthyroidism and diabetes mellitus may also occur.

**Hypersensitivity** has been described and may be exacerbated when aldesleukin is administered with other drugs; delayed reactions may occur with contrast media.

**CNS toxicity** is dose-related and generally reversible, but it may be associated with demyelination and may be irreversible in some patients.

[back to top](#)

**E - Dosing**

Refer to protocol by which patient is being treated. Patients must be managed as an inpatient in a tertiary care setting with full ICU facilities. Consider the following pre-medications (generally discontinued 12 hours after the last dose of aldesleukin):

- prophylactic antibiotics in patients with indwelling catheters
- Antipyretics
- H<sub>2</sub> antagonists
- Start meperidine and antidiarrheals early at the onset of symptoms.

Note: 1 MU = 1 million units = 1 million IU (international units) = 0.061 mg

**Adults:**

Optimal dosage and regimen for aldesleukin have not been established, but the following regimen has been approved by Health Canada.

**Intravenous:** 600,000 IU/kg (by a 15-minute infusion q8h; maximum 14 doses x 2)

Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
Dose	XXX	XXX	XXX	XXX	XX										XXX	XXX	XXX	XXX	XX

(X refers to a single dose)

The course may be repeated 7 weeks after hospital discharge, if tolerable and there is evidence of anti-tumour response.

**Dosage with Toxicity:**

Aldesleukin should be held for toxicity rather than reducing the dose to be given. Consult the product monograph for detailed management recommendations.

Dose interruption:

Body system		Hold; May Restart after Recovery	Discontinue permanently
Cardiovascular	Atrial fibrillation, SVT, bradycardia	X	
	Hypotension – pressors needed	X	
	EKG ischemia or myocarditis	X	
	Sustained VT, uncontrollable arrhythmia		X
	Angina, AMI, tamponade		X
Respiratory	O <sub>2</sub> saturation < 94% on room air or < 90% with 2 liters O <sub>2</sub>	X	
	Intubation for > 72hrs		X
CNS	Moderate-severe confusion, agitation, lethargy or somnolence	X	
	Coma, psychosis > 48 hrs		X
	Uncontrollable seizures		X
Infection	Grade 4, unstable	X	
Renal	Serum creatinine > 400 µmol/L Oliguria - < 10 mL/hour for 16-24 hours with ↑ creatinine	X	
	Dialysis required ≥ 72hrs		X
Hepatic	Hepatic failure**	X	
Skin	Bullous dermatitis	X	
GI	Ischemia, perforation		X
	Bleed not requiring surgery	X	
	Bleed requiring surgery		X

\*\*Abandon course. Do not start new course for at least 7 weeks after recovery.

Dosage in myelosuppression: No adjustment required

### **Dosage with Hepatic Impairment:**

Hold until recovery if any signs of hepatic failure are present. See table above.

**Dosage with Renal Impairment:**

Do not start treatment if creatinine > 130 µmol/L. See table above.

**Dosage in the elderly:**

Limited data in elderly patients. Monitor renal function closely as the elderly may have decreased renal function.

**Children:**

Safety and effectiveness in children under 18 years of age have not been established.

[back to top](#)

**F - Administration Guidelines**

- Each vial (1.3 mg) should be reconstituted with 1.2 mL of SWI.
- During reconstitution, USP should be directed at the side of the vial and the contents gently swirled to avoid excess foaming. Do not shake.
- Addition of BSWI or NS may promote aggregation and these solutions should not be used
- For IV administration, dilute further in 50 mL bag of D5W; infuse over 15 minutes
- Final concentration of drug should be 30 - 70µg/mL
- Drug delivery is more consistent when diluted in PVC container rather than non-PVC container
- In-line filter should not be used when administering aldesleukin
- Keep refrigerated; do not freeze. Avoid exposure to heat and light.

[back to top](#)

**G - Special Precautions****Contraindications:**

- known allergic reaction to aldesleukin, interleukin-2 or any components of the product
- abnormal thallium stress test

- abnormal pulmonary function tests
- organ allografts
- significant cardiac, pulmonary, renal, hepatic or CNS impairment
- concomitant use of cisplatin, vinblastine and dacarbazine as fatal tumour lysis syndrome has been reported

#### Other Warnings/Precautions:

- use with extreme caution in patients with known cardiac, pulmonary or seizure disorders (even with normal thallium and pulmonary function tests)
- use with extreme caution in patients with large fluid requirements (e.g., hypercalcemia)
- use with caution in patients with inflammatory or autoimmune disorders
- use with caution in combination with antihypertensives and psychotropic drugs as well as hepatotoxic, cardiotoxic or nephrotoxic drugs

#### Other Drug Properties:

- Carcinogenicity: Unknown

#### Pregnancy and Lactation:

- Embryotoxicity: Probable  
Aldesleukin is not recommended for use in pregnancy. Adequate contraception should be used by both sexes during treatment, and for at least **6 months** after the last dose.
- Excretion into breast milk: Unknown  
Breastfeeding is not recommended.

#### Other:

Patients should have baseline pulmonary function tests with arterial blood gases, thallium stress tests to exclude the presence of significant coronary artery disease, and brain imaging to exclude metastases. Patients who have had a nephrectomy are still eligible for treatment if they have serum creatinine levels within normal range. Pre-existing infections must be treated prior to starting aldesleukin.

[back to top](#)

### H - Interactions

AGENT	EFFECT	MECHANISM	MANAGEMENT
Psychotropics (narcotics, sedatives, tranquilizers, antiemetics)	Enhances CNS toxicities	Additive	Caution



Antihypertensives	Potentiate hypotension	Additive	Caution; monitor blood pressure
Radiographic iodinated contrast media	non-anaphylactic hypersensitivity	Unknown	Caution
Glucocorticoids	↓ anti-tumour effect of aldesleukin	Unknown	Avoid, but may be needed to manage side effects
cytotoxics	Potentiate myelosuppression, TLS	Additive	Caution
Nephrotoxic drugs (i.e. aminoglycosides, amphotericin B, methotrexate)	potentiate renal toxicity	Additive (possibly)	Caution
Hepatotoxic agents (e.g., methotrexate, asparaginase)	potentiate hepatotoxicity	Additive (possibly)	Caution
Cardiotoxic agents (e.g., doxorubicin)	Potentiate cardiotoxicity	Additive (possibly)	Caution
DTIC, cisplatin, interferon, tamoxifen	↑ hypersensitivity	Additive	Caution
Interferon	↑ hypersensitivity, cardiotoxicity and rhabdomyolysis, autoimmune disease	Unknown	Caution
Drug excreted via liver or kidney	↓ excretion	↓ renal, liver function	Caution

[back to top](#)

## I - Recommended Clinical Monitoring

### Recommended Clinical Monitoring

Monitor Type	Monitor Frequency
CBC	Baseline and frequent

Electrolytes, blood glucose	Baseline and frequent
Liver function tests	Baseline and frequent
Renal function tests, fluid intake and output	Baseline and frequent
Pulmonary function tests with arterial blood gases (FEV <sub>1</sub> > 2L or ≥ 75% of predicted for age)	Baseline
CT scan of brain to exclude CNS metastases	Baseline
Thallium stress tests to exclude significant coronary artery disease	Baseline
Frequent weight, vital signs, EKG etc. during treatment	
Chest x-ray	Baseline and regular
Clinical toxicity assessment for GI, hydration, cardiovascular, CNS, perfusion, autoimmune effects	regular

Grade toxicity using the current [NCI-CTCAE \(Common Terminology Criteria for Adverse Events\) version](#)

### **Suggested Clinical Monitoring**

Monitor Type	Monitor Frequency
<ul style="list-style-type: none"> <li>Repeat thallium testing if suspicion of ischemia or congestive cardiac failure</li> <li>Constant cardiac rhythm monitoring and hourly vital signs in patients with hypotension, especially bp &lt; 90 mmHg</li> </ul>	

[back to top](#)

## **J - Supplementary Public Funding**

### **New Drug Funding Program ([NDFP Website](#))**

- Aldesleukin (interleukin-2) - In-Transit Metastases from Melanoma

[back to top](#)

## **K - References**

McEvoy GK, editor. AHFS Drug Information 2009. Bethesda: American Society of Health-System Pharmacists, p. 915-24.

---

Proleukin® (aldesleukin/interleukin-2) [product monograph]. Dorval, Quebec: Novartis Pharmaceuticals Canada Inc.; Sept 20, 2012.

**June 2019** Updated emetic risk category

[back to top](#)

## L - Disclaimer

Refer to the [New Drug Funding Program](#) or [Ontario Public Drug Programs](#) websites for the most up-to-date public funding information.

*The information set out in the drug monographs, regimen monographs, appendices and symptom management information (for health professionals) contained in the Drug Formulary (the "Formulary") is intended for healthcare providers and is to be used for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or adverse effects of a particular drug, nor should it be construed to indicate that use of a particular drug is safe, appropriate or effective for a given condition. The information in the Formulary is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.*

*The format and content of the drug monographs, regimen monographs, appendices and symptom management information contained in the Formulary will change as they are reviewed and revised on a periodic basis. The date of last revision will be visible on each page of the monograph and regimen. Since standards of usage are constantly evolving, it is advised that the Formulary not be used as the sole source of information. It is strongly recommended that original references or product monograph be consulted prior to using a chemotherapy regimen for the first time.*

*Some Formulary documents, such as the medication information sheets, regimen information sheets and symptom management information (for patients), are intended for patients. Patients should always consult with their healthcare provider if they have questions regarding any information set out in the Formulary documents.*

*While care has been taken in the preparation of the information contained in the Formulary, such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability.*

*CCO and the Formulary's content providers shall have no liability, whether direct, indirect, consequential, contingent, special, or incidental, related to or arising from the information in the Formulary or its use thereof, whether based on breach of contract or tort (including negligence), and even if advised of the possibility thereof. Anyone using the*

---

---

*information in the Formulary does so at his or her own risk, and by using such information, agrees to indemnify CCO and its content providers from any and all liability, loss, damages, costs and expenses (including legal fees and expenses) arising from such person's use of the information in the Formulary.*

[back to top](#)