

# asciminib

**Pronunciation:** as-SIM-i-nib

**Other Name(s):** Scemblix™

**Appearance:** round tablets in various strengths and colours

This handout gives general information about this cancer medication.

You will learn:

- who to contact for help
- what the medication is
- how it is given
- what to expect while on medication



This handout was created by Ontario Health (Cancer Care Ontario) together with patients and their caregivers who have also gone through cancer treatment. It is meant to help support you through your cancer treatment and answer some of your questions.

**This information does not replace the advice of your health care team. Always talk to your health care team about your treatment.**

## Who do I contact if I have questions or need help?

My cancer health care provider is: \_\_\_\_\_

During the day I should contact: \_\_\_\_\_

Evenings, weekends and holidays: \_\_\_\_\_

## What is this treatment for?

For treating a type of blood cancer called chronic myeloid leukemia (CML)

## What should I do before I start this treatment?

Tell your health care team if you have or had significant medical condition(s), especially if you have / had:

- heart problems, including irregular heartbeat,
- serious infections, including hepatitis B
- pancreas problems,
- or any allergies

This drug contains a small amount of lactose. If you cannot tolerate lactose, talk to your health care team.



### Remember To:

- ✓ Tell your health care team about all of the other medications you are taking.
- ✓ Keep taking other medications that have been prescribed for you, unless you have been told not to by your health care team.

You will have a blood test to check for hepatitis B before starting treatment. See the [Hepatitis B and Cancer Medications](#) pamphlet for more information.

## How is this treatment given?

- Talk to your health care team about how and when to take your medication.
- Take the dose(s) at about the same time(s) each day.
- Swallow whole pill with a glass of water on an empty stomach, at least one hour before or two hours after food.
- Do not crush or chew the tablets.
- If you take asciminib **once daily** and miss a dose, take it if it is within 12 hours of the missed dose, otherwise skip and take your next dose as scheduled. Do not double the dose to make up for the forgotten one.
- If you take asciminib **twice daily** and miss a dose, take it if it is within 6 hours of the missed dose, otherwise skip and take your next dose as scheduled. Do not double the dose to make up for the forgotten one.
- If you vomit (throw up) after taking your medication, talk to your health care team about what to do.

**Warning:** If you take too much of this medication by accident, or if you think a child or a pet may have swallowed your medication, you must call the Ontario Poison Control Center right away at: 1-800-268-9017.

## Other medications you may be given with this treatment

### To Prevent Hepatitis B Flare Ups

If you have ever been infected with hepatitis B, there is a risk that this treatment can cause it to flare up (come back). Tell your health care team if you have had hepatitis B. You may need to take medication to prevent a hepatitis B flare-up.

**DO this while on treatment**

- ✓ DO check with your health care team before getting any vaccinations, surgery, dental work or other medical procedures.
- ✓ DO tell your health care team about any serious infections that you have now or have had in the past.
- ✓ DO protect your skin from the sun. Wear a long sleeved shirt, long pants and a hat. Apply sunscreen with UVA and UVB protection and an SPF of at least 30. Your skin may be more sensitive to the sun and you could develop a bad sunburn or rash more easily.

**DO NOT do this while on treatment**

- ✗ DO NOT take any other medications, such as vitamins, over-the-counter (non-prescription) drugs, or natural health products without checking with your health care team.
- ✗ DO NOT start any complementary or alternative therapies, such as acupuncture or homeopathic products, without checking with your health care team.
- ✗ DO NOT use tobacco products (such as smoking cigarettes or vaping) or drink alcohol while on treatment without talking to your health care team first. Smoking and drinking can make side effects worse and make your treatment not work as well.

## Will this treatment interact with other medications or natural health products?

Yes, this medication can interact with other medications, vitamins, foods and natural health products. Interactions can make this medication not work as well or cause severe side effects.

Tell your health care team about all of your:

- prescription and over-the-counter (non-prescription) medications
- other drugs and substances, such as cannabis/marijuana (medical or recreational)
- natural health products such as vitamins, herbal teas, homeopathic medicines, and other supplements, or traditional medicines

Check with your health care team before starting or stopping any of them.

If you take seizure medications (such as phenytoin), your health care team may monitor your blood levels closely and may change your dose.



### Talk to your health care team **BEFORE** taking or using these :

- Anti-inflammatory medications such as ibuprofen (Advil<sup>®</sup> or Motrin<sup>®</sup>), naproxen (Aleve<sup>®</sup>) or Aspirin<sup>®</sup>.
- Over-the-counter products such as dimenhydrinate (Gravol<sup>®</sup>)
- Natural health products such as St. John's Wort
- Traditional medicines
- Supplements such as vitamin C
- Grapefruit juice
- Alcoholic drinks
- Tobacco
- All other drugs, such as marijuana or cannabis (medical or recreational)

## What to do if you feel unwell, have pain, a headache or a fever

- ✓ **Always** check your temperature to see if you have a fever **before** taking any medications for fever or pain (such as acetaminophen (Tylenol®) or ibuprofen (Advil®)).
  - Fever can be a sign of infection that may need treatment right away.
  - If you take these medications before you check for fever, they may lower your temperature and you may not know you have an infection.

### How to check for fever:

Keep a digital (electronic) thermometer at home and take your temperature if you feel hot or unwell (for example, chills, headache, mild pain).

- You have a fever if your temperature taken **in your mouth (oral temperature)** is:
  - 38.3°C (100.9°F) or higher at any time

OR

  - 38.0°C (100.4°F) or higher for at least one hour.



### If you do have a fever:

- ✓ **Try to contact your health care team. If you are not able to talk to them for advice, you MUST get emergency medical help right away.**
- ✓ Ask your health care team for the [Fever](#) pamphlet for more information.

If you do not have a fever but have mild symptoms such as headache or mild pain:

- ✓ Ask your health care team about the right medication for you. **Acetaminophen (Tylenol®)** is a safe choice for most people.



**Talk to your health care team before you start taking** ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or ASA (Aspirin®), as they may increase your chance of bleeding or interact with your cancer treatment.



Talk to your health care team if you already take **low dose aspirin** for a medical condition (such as a heart problem). It may still be safe to take.

## How will this treatment affect sex, pregnancy and breastfeeding?

**Talk to your health care team about:**

- How this medication may affect your sexual health.
- How this medication may affect your ability to have a baby, if this applies to you.

This medication may harm an unborn baby. Tell your health care team if you or your partner are pregnant, become pregnant during treatment, or are breastfeeding.

- If there is **any** chance of pregnancy happening, you and your partner together must use **2 effective forms of birth control** at the same time until at least **1 week** after your last dose. Talk to your health care team about which birth control options are best for you.
- Do not breastfeed while on this medication and for at least **1 week** after your last dose.

## How to safely store and handle this medication

- Keep this medication in the original packaging at room temperature in a dry place, away from heat and light. Keep out of sight and reach of children and pets.
- Do not throw out any unused medications at home. Bring them to your pharmacy to be thrown away safely.

## How to safely touch oral anti-cancer medication

### If you are a patient:

- Wash your hands before and after touching your oral anti-cancer medication.
- Swallow each pill whole. Do not crush or chew your pills.

### If you are a caregiver:

- Wear nitrile or latex gloves when touching tablets, capsules or liquids.
- Wash your hands before putting on your gloves and after taking them off, even if your skin did not touch the oral anti-cancer medication.
- Throw out your gloves after each use. Do not re-use gloves.
- Do not touch oral anti-cancer medications if you are pregnant or breastfeeding.

## What to do if anti-cancer medication gets on your skin or in your eyes

### If medication gets on your skin:

- Wash your skin with a lot of soap and water.
- If your skin gets red or irritated, talk to your health care team.

### If medication gets in your eyes:

- Rinse your eyes with running water right away. Keep water flowing over your open eyes for at least 15 minutes.



## What are the side effects of this treatment?

The following table lists side effects that you may have when getting asciminib. The table is set up to list the most common side effects first and the least common last. It is unlikely that you will have all of the side effects listed and you may have some that are not listed.

Read over the side effect table so that you know what to look for and when to get help. Refer to this table if you experience any side effects while on asciminib.

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<p><b>Allergic reaction</b></p> <p>(May rarely be severe)</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>Hives, itchiness with or without rash, swollen lips, face or tongue, chest and throat tightness.</li> </ul> <p><b>What to do?</b></p> <ul style="list-style-type: none"> <li>Do not manage these symptoms on your own. Talk to your health care team right away if you have any of these symptoms.</li> <li>If you are unable to talk to the team for advice, you must get emergency medical help right away for severe symptoms and any problems with breathing.</li> </ul>	<p>Talk to your health care team right away if you have any of these symptoms.</p> <p>If you are unable to talk to the team for advice, you must get emergency medical help right away for severe symptoms and any problems with breathing.</p>
<p><b>Low neutrophils (white blood cells) in the blood (neutropenia)</b></p> <p>(May be severe)</p> <p>When neutrophils are low, you are at risk of getting an infection more easily. Ask your health care team for the <a href="#">Neutropenia (Low white blood cell count)</a> pamphlet for more information.</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>If you feel hot or unwell (for example if you have chills or a new cough), you <b>must</b> check your temperature to see if you have a fever.</li> <li>Do not take medications that treat a fever before you take your temperature (for example, Tylenol® (acetaminophen), or Advil® (ibuprofen)).</li> <li>Do not eat or drink anything hot or cold right before taking your temperature.</li> </ul>	<p>If you have a fever, try to contact your health care team. If you are unable to talk to the team for advice, you <b>MUST</b> get emergency medical help right away.</p>

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>You have a fever if your temperature taken in your mouth (oral temperature) is:</p> <ul style="list-style-type: none"> <li>• 38.3°C (100.9°F) or higher at any time</li> <li>OR</li> <li>• 38.0°C (100.4°F) or higher for at least one hour.</li> </ul> <p><b>What to do?</b></p> <p><b>If your health care team has told you that you have low neutrophils:</b></p> <ul style="list-style-type: none"> <li>• Wash your hands often to prevent infection.</li> <li>• Check with your health care team before getting any vaccines, surgeries, medical procedures or visiting your dentist.</li> <li>• Keep a digital thermometer at home so you can easily check for a fever.</li> </ul> <p><b>If you have a fever:</b></p> <p><b>If you have a fever, try to contact your health care team. If you are unable to talk to the team for advice, you must get emergency medical help right away.</b></p>	
<p><b>Low platelets in the blood</b></p> <p>(May be severe)</p> <p>When your platelets are low, you are at risk for bleeding and bruising. Ask your health care team for the <a href="#">Low Platelet Count</a> pamphlet for more information.</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Watch for signs of bleeding: <ul style="list-style-type: none"> <li>◦ bleeding from your gums</li> <li>◦ unusual or heavy nosebleeds</li> <li>◦ bruising easily or more than normal</li> <li>◦ black coloured stools (poo) or blood in your stools (poo)</li> <li>◦ coughing up red or brown coloured mucus</li> <li>◦ dizziness, constant headache or changes in your vision</li> <li>◦ heavy vaginal bleeding</li> </ul> </li> </ul>	<p>Talk to your health care team if you have any signs of bleeding. If you have bleeding that doesn't stop or is severe (very heavy), you <b>MUST</b> get emergency help right away.</p>

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<p><b>What to do?</b></p> <p><b>If your health care team has told you that you have low platelets:</b></p> <ul style="list-style-type: none"> <li>• Tell your pharmacist that your platelet count may be low before taking any prescriptions or over-the-counter medication.</li> <li>• Check with your healthcare team before you go to the dentist.</li> <li>• Take care of your mouth and use a soft toothbrush.</li> <li>• Try to prevent cuts and bruises.</li> <li>• Ask your health care team what activities are safe for you.</li> <li>• Your treatment may have to be delayed if you have low platelets. Your health care team may recommend a blood transfusion.</li> </ul> <p><b>If you have signs of bleeding:</b></p> <ul style="list-style-type: none"> <li>• If you have a small bleed, clean the area with soap and water or a saline (saltwater) rinse. Apply pressure for at least 10 minutes.</li> </ul> <p><b>If you have bleeding that does not stop or is severe (very heavy), you must get emergency medical help right away.</b></p>	
<p><b>Anemia (low red blood cells)</b></p> <p>(May be severe)</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• You may feel more tired or weaker than normal.</li> <li>• Pale skin and cold hands and feet.</li> <li>• You may feel short of breath, dizzy or lightheaded.</li> <li>• This may occur in days to weeks after your treatment starts.</li> </ul> <p><b>What to do?</b></p> <p><b>If your health care team has told you that you have anemia (low red blood cells):</b></p> <ul style="list-style-type: none"> <li>• Rest often and eat well.</li> <li>• Light exercise, such as walking may help.</li> <li>• You may need medication or a blood transfusion.</li> <li>• If it is very bad, your doctor may need to make changes to your treatment regimen.</li> </ul>	<p>Talk to your health care team if it does not improve or if it is severe.</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p><b>Higher than normal levels of pancreas enzymes in your blood (lipase, amylase)</b></p> <p>(May be severe)</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• These changes usually do not cause any symptoms.</li> <li>• In some cases, you may feel pain in the centre of your belly that may spread to your back.</li> </ul> <p><b>What to do?</b></p> <ul style="list-style-type: none"> <li>• Your health care team may do blood tests to watch your levels of enzymes in the pancreas.</li> </ul> <p><b>If you have any symptoms, get emergency medical help right away.</b></p>	<p>Talk to your health care team. If you are unable to talk to the team for advice, you must get emergency medical help right away.</p>
<p><b>Mild joint, muscle pain or cramps</b></p> <p><b>Headache</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• New pain in your muscles or joints, muscle cramps, or feeling achy.</li> <li>• Mild headache</li> </ul> <p><b>What to do?</b></p> <ul style="list-style-type: none"> <li>• Take pain medication (acetaminophen or opioids such as codeine, morphine, hydromorphone, oxycodone) as prescribed.</li> <li>• Read the above section: "<b>What to do if I feel unwell, have pain, a headache or a fever?</b>" before taking acetaminophen (Tylenol®), ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or Aspirin. These medications may hide an infection that needs treatment or they may increase your risk of bleeding.</li> <li>• Rest often and try light exercise (such as walking) as it may help.</li> </ul> <p>Ask your health care team for the <a href="#">Pain</a> pamphlet for more information.</p>	<p>Talk to your health care team if it does not improve or if it is severe.</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p><b>Fatigue</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>Feeling of tiredness or low energy that lasts a long time and does not go away with rest or sleep.</li> </ul> <p><b>What to do?</b></p> <ul style="list-style-type: none"> <li>Be active. Aim to get 30 minutes of moderate exercise (you are able to talk comfortably while exercising) on most days.</li> <li>Check with your health care team before starting any new exercise.</li> <li>Pace yourself, do not rush. Put off less important activities. Rest when you need to.</li> <li>Ask family or friends to help you with things like housework, shopping, and child or pet care.</li> <li>Eat well and drink at least 6 to 8 glasses of water or other liquids every day (unless your health care team has told you to drink more or less).</li> <li>Avoid driving or using machinery if you are feeling tired.</li> </ul> <p>Ask your health care team for the <a href="#">Fatigue</a> pamphlet for more information.</p>	<p>Talk to your health care team if it does not improve or if it is severe.</p>
<p><b>High blood pressure</b></p> <p>(May be severe)</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>There are usually no signs of high blood pressure.</li> <li>Rarely, you may have headaches, shortness of breath or nosebleeds.</li> </ul> <p><b>What to do?</b></p> <ul style="list-style-type: none"> <li>Check your blood pressure regularly.</li> <li>Your doctor may prescribe medication to treat high blood pressure.</li> </ul> <p><b>If you have a severe headache get emergency help right away as it may be a sign your blood pressure is too high.</b></p>	<p>Talk to your health care team if it does not improve or if it is severe.</p>
<p><b>Rash; dry, itchy skin</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>You may have cracked, rough, flaking or peeling areas of the skin.</li> <li>Your skin may look red and feel warm, like a sunburn.</li> <li>Your skin may itch, burn, sting or feel very tender when touched.</li> </ul>	<p>Talk to your health care team if it does not improve or if it is severe.</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p><b>What to do?</b></p> <p><b>To prevent and treat dry skin:</b></p> <ul style="list-style-type: none"> <li>• Use fragrance-free skin moisturizer.</li> <li>• Protect your skin from the sun and the cold.</li> <li>• Use sunscreen with UVA and UVB protection and a SPF of at least 30.</li> <li>• Avoid perfumed products and lotions that contain alcohol.</li> <li>• Drink 6 to 8 cups of non-alcoholic, non-caffeinated liquids each day, unless your health care team has told you to drink more or less.</li> </ul> <p><b>Rash may be severe in some rare cases and cause your skin to blister or peel. If this happens, get emergency medical help right away.</b></p>	
<p><b>Diarrhea</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Loose, watery, unformed stool (poo) that may happen days to weeks after you get your treatment.</li> </ul> <p><b>What to do?</b></p> <p><b>If you have diarrhea:</b></p> <ul style="list-style-type: none"> <li>• Take anti-diarrhea medication if your health care team prescribed it or told you to take it.</li> <li>• Do not eat foods or drinks with artificial sweetener (like chewing gum or 'diet' drinks), coffee and alcohol, until your diarrhea has stopped.</li> <li>• Eat many small meals and snacks instead of 2 or 3 large meals.</li> <li>• Drink at least 6 to 8 cups of liquids each day, unless your health care team has told you to drink more or less.</li> <li>• Talk to your health care team if you can't drink 6 to 8 cups of liquids each day when you have diarrhea. You may need to drink special liquids with salt and sugar, called Oral Rehydration Therapy.</li> <li>• Talk to your health care team if your diarrhea does not improve after 24 hours of taking diarrhea medication or if you have diarrhea more than 7 times in one day.</li> </ul> <p>Ask your health care team for the <a href="#">Diarrhea</a> pamphlet for more information.</p>	<p>Talk to your health care team if no improvement after 24 hours of taking diarrhea medication or if severe (more than 7 times in one day).</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p><b>Nausea and vomiting</b></p> <p>(Generally mild)</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Nausea is feeling like you need to throw up. You may also feel light-headed.</li> <li>• You may feel nausea within hours to days after your treatment.</li> </ul> <p><b>What to do?</b></p> <p><b>To help prevent nausea:</b></p> <ul style="list-style-type: none"> <li>• It is easier to prevent nausea than to treat it once it happens.</li> <li>• If you were given anti-nausea medication(s), take them as prescribed, even if you do not feel like throwing up.</li> <li>• Drink clear liquids and have small meals. Get fresh air and rest.</li> <li>• Do not eat spicy, fried foods or foods with a strong smell.</li> <li>• Limit caffeine (like coffee, tea) and avoid alcohol.</li> </ul> <p><b>If you have nausea or vomiting:</b></p> <ul style="list-style-type: none"> <li>• Take your rescue (as-needed) anti-nausea medication(s) as prescribed.</li> <li>• Ask your health care team for the <a href="#">Nausea &amp; Vomiting</a> pamphlet for more information.</li> <li>• Talk to your health care team if: <ul style="list-style-type: none"> <li>◦ nausea lasts more than 48 hours</li> <li>◦ vomiting lasts more than 24 hours or if it is severe</li> </ul> </li> </ul>	<p>Contact your healthcare team if nausea lasts more than 48 hours or vomiting lasts more than 24 hours.</p>

**Other rare, but serious side effects are possible with this treatment.**

If you have **any** of the following, talk to your cancer health care team or get emergency medical help right away:

- irregular heartbeat, shortness of breath, or fainting spells
- swelling in your legs, ankles and belly
- pain, swelling and hardening of a vein
- sharp pain in the centre or left side of the chest that may spread to your arm
- new cough, difficulty breathing or coughing up blood
- feeling confused, having a sudden loss of vision or trouble speaking or using your arms or legs

For more information on how to manage your symptoms ask your health care provider, or visit:

<https://www.cancercareontario.ca/symptoms>.

## Notes

---

---

---

---

---

### July 2025 New drug information sheet

*The information set out in the medication information sheets, regimen information sheets, and symptom management information (for patients) contained in the Drug Formulary (the "Formulary") is intended to be used by health professionals and patients for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or side effects of a certain drug, nor should it be used to indicate that use of a particular drug is safe, appropriate or effective for a given condition.*

*A patient should always consult a healthcare provider if he/she has any questions regarding the information set out in the Formulary. The information in the Formulary is not intended to act as or replace medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.*