

Drug Monograph

[Drug Name](#) | [Mechanism of Action and Pharmacokinetics](#) | [Indications and Status](#) | [Adverse Effects](#) | [Dosing](#) | [Administration Guidelines](#) | [Special Precautions](#) | [Interactions](#) | [Recommended Clinical Monitoring](#) | [Supplementary Public Funding](#) | [References](#) | [Disclaimer](#)

A - Drug Name

chlorambucil

COMMON TRADE NAME(S): Leukeran®

[back to top](#)

B - Mechanism of Action and Pharmacokinetics

Chlorambucil was synthesized in 1953. It is an aromatic derivative of mechlorethamine, is closely related in structure to melphalan and is a bifunctional alkylating agent. DNA alkylation by the reactive ethylenimonium radical results in breaks in the DNA molecule as well as cross-linking of the twin strands, thus interfering with DNA replication and transcription of RNA. Like other alkylators, chlorambucil is cell cycle phase-nonspecific. The immunosuppressive activity of chlorambucil is due to its suppression of lymphocytes.

Absorption	Bioavailability	oral: Yes, bioavailability 70-80%. Food decreases bioavailability by 10-20%.
------------	-----------------	--

Distribution	Rapidly cleared from plasma; tissue distribution fairly homogeneous; crosses placenta; found in ascitic fluid.
--------------	--

Cross blood brain barrier?	no information found, but can cause seizures in high doses.
----------------------------	---

Volume of distribution	0.14-0.24 L/kg
------------------------	----------------

PPB	99% (albumin)
-----	---------------

Metabolism	Extensively metabolized in liver by the hepatic microsomal enzyme oxidation system
------------	--

Active metabolites	phenylacetic acid mustard.
--------------------	----------------------------

	Inactive metabolites	Yes
Elimination	Extremely low urinary excretion. Chlorambucil and its metabolites are probably not dialysable.	
Urine	15-60% in 24 hours; less than 1% as intact drug / active metabolite.	
Half-life	1.5 hours	

[back to top](#)

C - Indications and Status

Health Canada Approvals:

- Monotherapy in chronic lymphocytic leukemia
- Monotherapy or combination therapy in:
 - Non-Hodgkin's lymphoma (e.g. follicular, MALT, mantle cell lymphoma, Waldenstrom's macroglobulinemia)
 - Hodgkin's lymphoma (palliative)

[back to top](#)

D - Adverse Effects

Emetogenic Potential: Minimal – No routine prophylaxis; PRN recommended

Extravasation Potential: Not applicable

ORGAN SITE	SIDE EFFECT* (%)	ONSET**
Dermatological	Rash (may be severe)	E
Gastrointestinal	Diarrhea (infrequent)	I
	Mucositis (infrequent)	E
	Nausea, vomiting (infrequent)	I
Hematological	Immunosuppression	E
	Myelosuppression ± infection, bleeding (may be	E

severe)		
Hepatobiliary	Hepatotoxicity	D
Hypersensitivity	Drug reaction	I
Metabolic / Endocrine	Tumor lysis syndrome	I
Neoplastic	Leukemia (secondary) (or other malignancies)	L
Nervous System	Neurotoxicity	E
	Peripheral neuropathy	L
	Seizure (Rare, with high doses, prior history)	E
Reproductive and breast disorders	Infertility	L
Respiratory	Pneumonitis	E
Urinary	Cystitis (sterile)	E

* "Incidence" may refer to an absolute value or the higher value from a reported range.
 "Rare" may refer to events with < 1% incidence, reported in post-marketing, phase 1 studies,
 isolated data or anecdotal reports.

** I = *immediate* (onset in hours to days) E = *early* (days to weeks)
 D = *delayed* (weeks to months) L = *late* (months to years)

Myelosuppression is more severe following continuous administration than intermittent high dose chlorambucil and is dose-related. Prolonged therapy or excessive doses may result in pancytopenia or irreversible bone marrow damage.

Hyperuricemia during periods of active cell lysis, which is caused by cytotoxic chemotherapy of highly proliferative tumours of massive burden (e.g. some leukemias and lymphomas), can be minimized with allopurinol and hydration. In hospitalized patients, the urine may be alkalinized, by addition of sodium bicarbonate to the IV fluids, if tumour lysis is expected.

Pulmonary toxicity similar to other alkylating agents can occur, including interstitial pulmonary fibrosis or pneumonitis. Signs and symptoms are dyspnea, dry cough, fever, rales and tachypnea developing over a 1-2 month period. It is usually associated with prolonged therapy (6-24 months) and related to a total dose of >2 g. Partial recovery can occur within several weeks after discontinuing therapy and administration of steroids. In other patients, pulmonary complications progress and some deaths have occurred.

Children with nephrotic syndrome and patients receiving high pulse doses or with a prior history of seizures may have an increased risk of **seizures**, and should be closely monitored. Rare, focal and/or generalized seizures have also occurred in patients receiving therapeutic daily doses.

Rarely, **skin rash** may occur and result in Stevens-Johnson Syndrome, toxic epidermal necrolysis or erythema multiforme. Chlorambucil should be discontinued immediately if skin reactions occur. Patients with a history of rash with other alkylating agents may have an increased risk of rash.

[back to top](#)

E - Dosing

Refer to protocol by which patient is being treated. Numerous dosing schedules exist and depend on disease, response and concomitant therapy. Guidelines for dosing also include consideration of white blood cell count. Dosage may be reduced and/or delayed in patients with bone marrow depression due to cytotoxic/radiation therapy.

CLL patients with pancytopenia should receive prednisone initially, and chlorambucil should only be started with some evidence of recovery – the initial dose of chlorambucil when started should not exceed 0.1 mg/kg/day.

Adults:

Continuous regimens:

- Induction: 0.1-0.2 mg/kg/day PO (CLL - until white cell count $10 \times 10^9/L$; for 4-8 weeks for NHL)
- Maintenance: After 4-week “rest”: 0.03-0.1 mg/kg/day PO adjusted to tolerance

Intermittent regimens:

- q28d: 6 mg/m²/d PO for 7-14 days

Dosage with Toxicity:

Modify according to protocol by which patient is being treated; if no guidelines available, refer to Appendix 6 " Dosage Modification for Hematologic and Non-Hematologic Toxicities."

Toxicity and grade	Action	Dose when restart
Severe myelosuppression/bone marrow failure	Discontinue	N/A
Grade 3 hematologic	Hold*	↓ by 33%
Grade 4 hematologic, febrile neutropenia or thrombocytopenic bleeding	Hold*	↓ by 33% or Discontinue
Grade 3 non-hematologic	Hold*	↓ by 33%

Skin or pulmonary toxicity or Grade 4 non-hematologic	Discontinue	N/A
--	-------------	-----

*Before retreatment, major organ toxicities should recover to ≤ Grade 2, platelets $\geq 100 \times 10^9/L$, and ANC $\geq 1.5 \times 10^9/L$.

Dosage with Hepatic Impairment:

Adjustment required with severe hepatic impairment; no details found.

Dosage with Renal Impairment:

No adjustment required, but monitor carefully as at increased risk of myelosuppression.

The following dosage adjustments have been used by some clinicians:

Creatinine Clearance (mL/min)	% Dose
10-50	75%
<10	50%

Dosage in the elderly:

No overall differences in safety or effectiveness were observed between younger patients and patients ≥ 65 years. Use with caution due to possible decreases in hepatic, renal, or cardiac function.

Children:

Safety and efficacy have not been established.

[back to top](#)

F - Administration Guidelines

- Oral self-administration; drug available by outpatient prescription.
- Keep drug refrigerated; do not freeze

[back to top](#)

G - Special Precautions

Other:

Chlorambucil is **contraindicated** in patients who are resistant to the drug or who have developed hypersensitivity to it; there may be cross-sensitivity with other alkylating agents (especially rash).

Should not be used within 4 weeks of a full course of radiation therapy or chemotherapy.

Administer with caution if bone marrow is severely depressed and in patients with seizure disorders. Avoid live vaccines.

Chlorambucil has been shown to have **teratogenic, mutagenic** and **carcinogenic** properties in experimental models and should not be used in **pregnancy**. Adequate contraception must be used by both sexes, during treatment and for at least 6 months after chlorambucil cessation. Chlorambucil is known to cross the placenta. **Breast feeding** is not recommended because of the potential secretion of chlorambucil into breast milk. Secondary malignancies have been reported.

Both reversible and permanent **sterility** have been observed in both sexes. Children receiving chlorambucil before puberty generally have a normal progression of puberty. In males, however, testicular atrophy may occur and persist.

[back to top](#)

H - Interactions

AGENT	EFFECT	MECHANISM	MANAGEMENT
Phenylbutazone	↑ the toxicity of chlorambucil	Unknown	Modify chlorambucil doses
Succinylcholine	prolonged apnea	inhibition of serum cholinesterase	decrease dose of succinylcholine
Live vaccines	systemic viral infection risk	immunosuppression	avoid / caution

[back to top](#)

I - Recommended Clinical Monitoring

Recommended Clinical Monitoring

Monitor Type	Monitor Frequency
--------------	-------------------

CBC	Weekly or twice weekly
Toxicity assessment (seizures, GI, hypersensitivity, TLS, infection, bleeding, pulmonary)	Routinely

Grade toxicity using the current [NCI-CTCAE \(Common Terminology Criteria for Adverse Events\) version](#)

Suggested Clinical Monitoring

Monitor Type	Monitor Frequency
Liver function tests	Periodic

[back to top](#)

J - Supplementary Public Funding

ODB - General Benefit ([ODB Formulary](#))

- chlorambucil ()

[back to top](#)

K - References

Aronoff GR, Bennett WM, Berns JS, et al, Drug Prescribing in Renal Failure: Dosing Guidelines for Adults and Children, 5th ed. Philadelphia, PA: American College of Physicians; 2007, p 98.

Bauwens D, Maerevoet M, Michaux L, et al. Activity and safety of combined rituximab with chlorambucil in patients with mantle cell lymphoma. British Journal of Haematology 2005; 131: 338–40.

Cancer Drug Manual (the Manual), 1994, British Columbia Cancer Agency (BCCA).

Chlorambucil: e-Drugdex, Micromedex Healthcare Series.

McEvoy GK, editor. AHFS Drug Information 2009. Bethesda: American Society of Health-System Pharmacists, p. 987-90.

Martinelli G, Laszlo D, Bertolini F, et al. Chlorambucil in combination with induction and maintenance rituximab is feasible and active in indolent non-Hodgkin's lymphoma. British Journal of Haematology 2003; 123: 271–7.

Radford JA, Rohatiner AZS, Ryder WDJ, et al. ChIVPP/EVA Hybrid Versus the Weekly VAPEC-B Regimen for Previously Untreated Hodgkin's Disease. J Clin Oncol 2002; 20: 2988-2994.

Product Monograph: Leukeran® (chlorambucil). Triton Pharma Inc. Canada, November 29, 2010.

Product Monograph: Leukeran® (chlorambucil). GlaxoSmithKline Inc. USA, October 2011.

June 2019 Updated emetic risk category

[back to top](#)

L - Disclaimer

Refer to the [New Drug Funding Program](#) or [Ontario Public Drug Programs](#) websites for the most up-to-date public funding information.

The information set out in the drug monographs, regimen monographs, appendices and symptom management information (for health professionals) contained in the Drug Formulary (the "Formulary") is intended for healthcare providers and is to be used for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or adverse effects of a particular drug, nor should it be construed to indicate that use of a particular drug is safe, appropriate or effective for a given condition. The information in the Formulary is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.

The format and content of the drug monographs, regimen monographs, appendices and symptom management information contained in the Formulary will change as they are reviewed and revised on a periodic basis. The date of last revision will be visible on each page of the monograph and regimen. Since standards of usage are constantly evolving, it is advised that the Formulary not be used as the sole source of information. It is strongly recommended that original references or product monograph be consulted prior to using a chemotherapy regimen for the first time.

Some Formulary documents, such as the medication information sheets, regimen information sheets and symptom management information (for patients), are intended for patients. Patients should always consult with their healthcare provider if they have questions regarding any information set out in the Formulary documents.

While care has been taken in the preparation of the information contained in the Formulary, such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability.

CCO and the Formulary's content providers shall have no liability, whether direct, indirect, consequential, contingent,

special, or incidental, related to or arising from the information in the Formulary or its use thereof, whether based on breach of contract or tort (including negligence), and even if advised of the possibility thereof. Anyone using the information in the Formulary does so at his or her own risk, and by using such information, agrees to indemnify CCO and its content providers from any and all liability, loss, damages, costs and expenses (including legal fees and expenses) arising from such person's use of the information in the Formulary.

[back to top](#)