

## Medication Information Sheet

**daSATinib** (da-SAT-in-ib)

*This document provides general information about your medication. It does not replace the advice of your health care professional. Always discuss your therapy with your health care professional and refer to the package insert for more details.*

**Other Name:** Sprycel®

**Appearance:** tablet In various strengths

**What is this medication for?**

- For treating certain types of blood cancers, such as chronic myeloid leukemia (CML) or acute lymphoblastic leukemia (ALL).

**What should I do before I have this medication?**

Tell your health care team if you have or had significant medical condition(s), especially if you have / had:

- liver disease (including hepatitis)
- heart problems (including irregular heartbeat) or
- any allergies
- Dasatinib tablets contains a small amount of lactose. If you cannot have lactose, talk to your healthcare team.

**Remember to:**

- Tell your health care team about all of the other medications you are taking.
- Keep taking other medications that have been prescribed for you, unless you have been told not to by your health care team.

**How will this medication affect sex, pregnancy and breastfeeding?**

**Talk to your health care team about:**

- How this medication may affect your sexual health.
- Changes to your menstrual cycle (periods), if this applies to you.

- How this medication may affect your ability to have a baby, if this applies to you.

This medication may harm an unborn baby. Tell your health care team if you or your partner are pregnant, become pregnant during treatment, or are breastfeeding.

- If there is **any** chance of pregnancy happening, you and your partner together must use **2 effective forms of birth control** at the same time until **6 months** after your last dose (general recommendation). Talk to your health care team about which birth control options are best for you.
- Do not breastfeed while using this medication.

## How is this medication given?

- Dasatinib is usually taken once a day by mouth; take the dose at about the same time each day.
- Swallow whole with a glass of water, with or without food.
- Do not crush or chew the tablets.
- If you miss a dose, skip this and take your next dose as scheduled. Do not double the dose to make up for the forgotten one.
- If you vomit (throw up) after taking your medication, talk to your health care team about what to do.
- If you take too much of your medication by accident, or if you think a child or a pet may have swallowed your medication, you must call the Ontario Poison Control Center right away at: 1-800-268-9017

## To Prevent Hepatitis B Flare Ups

If you have ever been infected with hepatitis B, there is a risk that this treatment can cause it to flare up (come back). Tell your health care team if you have had hepatitis B. You may need to take medication to prevent a hepatitis B flare-up.

## What else do I need to know while on this medication?

### Will this medication interact with other medications or natural health products?

- This medication can interact with other medications, vitamins, foods and natural health products. Interactions can make the treatment not work as well or cause severe side effects.
- Tell your health care team about all of your:

- prescription and over-the-counter (non-prescription) medications and all other drugs, such as marijuana (medical or recreational)
- natural health products such as vitamins, herbal teas, homeopathic medicines, and other supplements
- Check with your health care team before starting or stopping any of them.
- If you take seizure medications (such as phenytoin), your health care team may monitor your blood levels closely and may change your dose.
- If you are taking a blood thinner (such as warfarin), your health care team may need extra blood tests and may change your dose.
- Antacids should be avoided; if required, it should be taken at least 2 hours before or 2 hours after taking dasatinib.

### What should I do if I feel unwell, have pain, a headache or a fever?

- **Always** check your temperature to see if you have a fever **before** taking any medications for fever or pain (such as acetaminophen (Tylenol®) or ibuprofen (Advil®)).
  - Fever can be a sign of infection that may need treatment right away.
  - If you take these medications before you check for fever, they may lower your temperature and you may not know you have an infection.

### How to check for fever:

Keep a digital (electronic) thermometer at home and take your temperature if you feel hot or unwell (for example, chills, headache, mild pain).

- You have a fever if your temperature taken in your mouth (oral temperature) is:
  - 38.3°C (100.9°F) or higher at any time
  - OR
  - 38.0°C (100.4°F) or higher for at least one hour.

### If you do have a fever:

- **Try to contact your health care team. If you are not able to talk to them for advice, you MUST get emergency medical help right away.**
- Ask your health care team for the [Fever](#) pamphlet for more information.

**If you do not have a fever** but have mild symptoms such as headache or mild pain:

- Ask your health care team about the right medication for you. **Acetaminophen (Tylenol®)** is a safe choice for most people.
- **Talk to your health care team before you start taking** Ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or ASA (Aspirin®), as they may increase your chance of bleeding or interact with your cancer treatment.
- Talk to your health care team if you already take **low dose aspirin** for a medical condition (such as a heart problem). It may still be safe to take.

**What to DO while on this medication:**

- DO check with your health care team before getting any vaccinations, surgery, dental work or other medical procedures.

**What NOT to DO while on this medication:**

- DO NOT smoke or drink alcohol while on treatment without talking to your health care team first. Smoking and drinking can make side effects worse and make your treatment not work as well.
- DO NOT eat or drink grapefruit, starfruit, Seville oranges or their juices (or products that contain these) while taking this drug. They may increase the amount of drug in your blood and increase side effects.

## How should I safely store this medication?

- Keep this medication in the original packaging at room temperature in a dry place, away from heat and light. Keep out of sight and reach of children and pets.
- Do not throw out any unused medications at home. Bring them to your pharmacy to be thrown away safely.
- **How to safely touch oral anti-cancer medications**

**If you are a patient:**

- ◊ Wash your hands before and after touching your oral anti-cancer medication.
- ◊ Swallow each pill whole. Do not crush or chew your pills.

**If you are a caregiver:**

- ◊ Wear nitrile or latex gloves when touching tablets, capsules or liquids.
- ◊ Wash your hands before putting on your gloves and after taking them off, even if your skin did not touch the oral anti-cancer medication.
- ◊ Throw out your gloves after each use. Do not re-use gloves.
- ◊ Do not touch oral anti-cancer medications if you are pregnant or breastfeeding.

- **What to do if oral anti-cancer medication gets on your skin or in your eyes**

**If medication gets on your skin:**

- ◊ Wash your skin with a lot of soap and water.
- ◊ If your skin gets red or irritated, talk to your health care team.

**If medication gets in your eyes:**

- ◊ Rinse your eyes with running water right away. Keep water flowing over your open eyes for at least 15 minutes.

## What are the side effects of this medication?

The following table lists side effects that you may have when getting dasatinib. The table is set up to list the most common side effects first and the least common last. It is unlikely that you will have all of the side effects listed and you may have some that are not listed.

Read over the side effect table so that you know what to look for and when to get help. Refer to this table if you experience any side effects while on dasatinib.

<b>Common Side Effects (25 to 49 out of 100 people)</b>	
<b>Side effects and what to do</b>	<b>When to contact health care team</b>
<b>Mild joint, muscle pain or cramps</b>  <b>What to look for?</b> <ul style="list-style-type: none"> <li>• Mild headache</li> <li>• New pain in your muscles or joints, muscle cramps, or feeling achy.</li> </ul> <b>What to do?</b>	Talk to your health care team if it does not improve or if it is severe

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<ul style="list-style-type: none"> <li>Take pain medication (acetaminophen or opioids such as codeine, morphine, hydromorphone, oxycodone) as prescribed.</li> <li>Read the above section: <b>"What should I do if I feel unwell, have pain, a headache or a fever?"</b> before taking acetaminophen (Tylenol®), ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or Aspirin. These medications may hide an infection that needs treatment or they may increase your risk of bleeding.</li> <li>Rest often and try light exercise (such as walking) as it may help.</li> </ul> <p>Ask your health care team for the <a href="#">Pain</a> pamphlet for more information.</p>	
<p><b>Low neutrophils (white blood cells) in the blood (neutropenia)</b></p> <p>(May be severe)</p> <p>When neutrophils are low, you are at risk of getting an infection more easily. Ask your health care team for the <a href="#">Neutropenia (Low white blood cell count)</a> pamphlet for more information.</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>If you feel hot or unwell (for example if you have chills or a new cough), you <b>must</b> check your temperature to see if you have a fever.</li> <li>Do not take medications that treat a fever before you take your temperature (for example, Tylenol®, acetaminophen, Advil® or ibuprofen).</li> <li>Do not eat or drink anything hot or cold right before taking your temperature.</li> </ul> <p>You have a fever if your temperature taken in your mouth (oral temperature) is:</p> <ul style="list-style-type: none"> <li>38.3°C (100.9°F) or higher at any time</li> <li>OR</li> <li>38.0°C (100.4°F) or higher for at least one hour.</li> </ul> <p><b>What to do?</b></p> <p><b>If your health care team has told you that you have low neutrophils:</b></p> <ul style="list-style-type: none"> <li>Wash your hands often to prevent infection.</li> <li>Check with your health care team before getting any vaccines, surgeries, medical procedures or visiting your dentist.</li> <li>Keep a digital thermometer at home so you can easily check for a fever.</li> </ul> <p><b>If you have a fever:</b></p>	<p>If you have a fever, try to contact your health care team. If you are unable to talk to the team for advice, you <b>MUST</b> get emergency medical help right away.</p>

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<b>If you have a fever, try to contact your health care team. If you are unable to talk to the team for advice, you must get emergency medical help right away.</b>	
<p><b>Low platelets in the blood</b></p> <p>(May be severe)</p> <p>When your platelets are low, you are at risk for bleeding and bruising. Ask your health care team for the <a href="#">Low Platelet Count</a> pamphlet for more information.</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Watch for signs of bleeding: <ul style="list-style-type: none"> <li>◦ bleeding from your gums</li> <li>◦ unusual or heavy nosebleeds</li> <li>◦ bruising easily or more than normal</li> <li>◦ black coloured stools (poo) or blood in your stools (poo)</li> <li>◦ coughing up red or brown coloured mucus</li> <li>◦ dizziness, constant headache or changes in your vision</li> <li>◦ heavy vaginal bleeding</li> <li>◦ red or pink coloured urine (pee)</li> </ul> </li> </ul> <p><b>What to do?</b></p> <p><b>If your health care team has told you that you have low platelets:</b></p> <ul style="list-style-type: none"> <li>• Tell your pharmacist that your platelet count may be low before taking any prescriptions or over-the-counter medication.</li> <li>• Check with your healthcare team before you go to the dentist.</li> <li>• Take care of your mouth and use a soft toothbrush.</li> <li>• Try to prevent cuts and bruises.</li> <li>• Ask your health care team what activities are safe for you.</li> <li>• Your treatment may have to be delayed if you have low platelets. Your health care team may recommend a blood transfusion.</li> </ul> <p><b>If you have signs of bleeding:</b></p> <ul style="list-style-type: none"> <li>• If you have a small bleed, clean the area with soap and water or a saline (saltwater) rinse. Apply pressure for at least 10 minutes.</li> </ul> <p><b>If you have bleeding that does not stop or is severe (very heavy), you must get emergency medical help right away.</b></p>	<p>Talk to your health care team if you have any signs of bleeding. If you have bleeding that doesn't stop or is severe (very heavy), you <b>MUST</b> get emergency help right away.</p>

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<p><b>Diarrhea</b></p> <p>(May be severe)</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>Loose, watery, unformed stool (poo) that may happen days to weeks after you get your treatment.</li> </ul> <p><b>What to do?</b></p> <p><b>If you have diarrhea:</b></p> <ul style="list-style-type: none"> <li>Take anti-diarrhea medication if your health care team prescribed it or told you to take it.</li> <li>Do not eat foods or drinks with artificial sweetener (like chewing gum or 'diet' drinks), coffee and alcohol.</li> <li>Eat many small meals and snacks instead of 2 or 3 large meals.</li> <li>Drink at least 6 to 8 cups of liquids each day, unless your health care team has told you to drink more or less.</li> <li>Talk to your health care team if you can't drink 6 to 8 cups of liquids each day when you have diarrhea. You may need to drink special liquids with salt and sugar, called Oral Rehydration Therapy.</li> <li>Talk to your health care team if your diarrhea does not improve after 24 hours of taking diarrhea medication or if you have diarrhea more than 7 times in one day.</li> </ul> <p>Ask your health care team for the <a href="#">Diarrhea</a> pamphlet for more information.</p>	<p>Talk to your health care team if no improvement after 24 hours of taking diarrhea medication or if severe (more than 7 times in one day)</p>
<p><b>Fatigue</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>Feeling of tiredness or low energy that lasts a long time and does not go away with rest or sleep.</li> </ul> <p><b>What to do?</b></p> <ul style="list-style-type: none"> <li>Be active. Aim to get 30 minutes of moderate exercise (you are able to talk comfortably while exercising) on most days.</li> <li>Check with your health care team before starting any new exercise.</li> <li>Pace yourself, do not rush. Put off less important activities. Rest when you need to.</li> <li>Ask family or friends to help you with things like housework, shopping, and child</li> </ul>	<p>Talk to your health care team if it does not improve or if it is severe</p>



Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>or pet care.</p> <ul style="list-style-type: none"> <li>• Eat well and drink at least 6 to 8 glasses of water or other liquids every day (unless your health care team has told you to drink more or less).</li> <li>• Avoid driving or using machinery if you are feeling tired.</li> </ul> <p>Ask your health care team for the <a href="#">Fatigue</a> pamphlet for more information.</p>	
<p><b>Cough and feeling short of breath</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• You may have a cough and feel short of breath.</li> <li>• Symptoms that commonly occur with a cough are: <ul style="list-style-type: none"> <li>◦ wheezing or a whistling breathing</li> <li>◦ runny nose</li> <li>◦ sore throat</li> <li>◦ heartburn</li> <li>◦ weight loss</li> <li>◦ fever and chills</li> </ul> </li> <li>• Rarely this may be severe with chest pain, trouble breathing or coughing up blood.</li> </ul> <p><b>What to do?</b></p> <ul style="list-style-type: none"> <li>• Check your temperature to see if you have a fever. Read the above section <b>"What should I do if I feel unwell, have pain, a headache or a fever?"</b>.</li> <li>• If you have a fever, try to talk to your health care team. <b>If you are not able to talk to them for advice, you MUST get emergency medical help right away.</b></li> <li>• <b>If you have a severe cough with chest pain, trouble breathing or you are coughing up blood, get medical help right away.</b></li> </ul>	<p>Talk to your health care team. If you are not able to talk to your health care team for advice, and you have a fever or severe symptoms, you <b>MUST</b> get emergency medical help right away</p>
<p><b>Rash; dry, itchy skin</b></p> <p>(May be severe)</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• You may have cracked, rough, flaking or peeling areas of the skin.</li> <li>• Your skin may look red and feel warm, like a sunburn.</li> <li>• Your skin may itch, burn, sting or feel very tender when touched.</li> </ul> <p><b>What to do?</b></p> <p><b>To prevent and treat dry skin:</b></p>	<p>Talk to your health care team if it does not improve or if it is severe</p>

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<ul style="list-style-type: none"> <li>• Use fragrance-free skin moisturizer.</li> <li>• Protect your skin from the sun and the cold.</li> <li>• Use sunscreen with UVA and UVB protection and a SPF of at least 30.</li> <li>• Avoid perfumed products and lotions that contain alcohol.</li> <li>• Drink 6 to 8 cups of non-alcoholic, non-caffeinated liquids each day, unless your health care team has told you to drink more or less.</li> </ul> <p><b>Rash may be severe in some rare cases and cause your skin to blister or peel. It may also cause "bulls-eye" markings on your skin. If any of these symptoms happen, get emergency medical help right away.</b></p>	
<p><b>Swelling</b></p> <p>(May be severe)</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• You may have mild swelling or puffiness in your arms and/or legs. Rarely, this may be severe.</li> </ul> <p><b>What to do?</b></p> <p><b>To help prevent swelling:</b></p> <ul style="list-style-type: none"> <li>• Eat a low-salt diet.</li> </ul> <p><b>If you have swelling:</b></p> <ul style="list-style-type: none"> <li>• Wear loose-fitting clothing.</li> <li>• For swollen legs or feet, keep your feet up when sitting.</li> </ul>	<p>Talk to your health care team if it does not improve or if it is severe</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p><b>Nausea and vomiting</b></p> <p>(Generally mild)</p> <p><b>What to look for?</b></p>	<p>Talk to your healthcare team if nausea lasts more than 48 hours or vomiting lasts</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<ul style="list-style-type: none"> <li>• Nausea is feeling like you need to throw up. You may also feel light-headed.</li> <li>• You may feel nausea within hours to days after your treatment.</li> </ul> <p><b>What to do?</b></p> <p><b>To help prevent nausea:</b></p> <ul style="list-style-type: none"> <li>• It is easier to prevent nausea than to treat it once it happens.</li> <li>• If you were given anti-nausea medication(s), take them as prescribed, even if you do not feel like throwing up.</li> <li>• Drink clear liquids and have small meals. Get fresh air and rest.</li> <li>• Do not eat spicy, fried foods or foods with a strong smell.</li> <li>• Limit caffeine (like coffee, tea) and avoid alcohol.</li> </ul> <p><b>If you have nausea or vomiting:</b></p> <ul style="list-style-type: none"> <li>• Take your rescue (as-needed) anti-nausea medication(s) as prescribed.</li> <li>• Ask your health care team for the <a href="#">Nausea &amp; Vomiting</a> pamphlet for more information.</li> <li>• Talk to your health care team if:             <ul style="list-style-type: none"> <li>◦ nausea lasts more than 48 hours</li> <li>◦ vomiting lasts more than 24 hours or if it is severe</li> </ul> </li> </ul>	more than 24 hours or if it is severe
<p><b>Constipation</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Having bowel movements (going poo) less often than normal.</li> <li>• Small hard stools (poo) that look like pellets.</li> <li>• The need to push hard and strain to have any stool (poo) come out.</li> <li>• Stomach ache or cramps.</li> <li>• A bloated belly, feeling of fullness, or discomfort.</li> <li>• Leaking of watery stools (poo).</li> <li>• Lots of gas or burping.</li> <li>• Nausea or vomiting.</li> </ul> <p><b>What to do?</b></p> <p><b>To help prevent constipation:</b></p>	Talk to your health care team if it does not improve or if it is severe

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<ul style="list-style-type: none"> <li>• Try to eat more fiber rich foods like fruits with skin, leafy greens and whole grains.</li> <li>• Drink at least 6 to 8 cups of liquids each day unless your health care team has told you to drink more or less.</li> <li>• Be Active. Exercise can help to keep you regular.</li> <li>• If you take opioid pain medication, ask your health care team if eating more fibre is right for you.</li> </ul> <p><b>To help treat constipation:</b></p> <ul style="list-style-type: none"> <li>• If you have not had a bowel movement in 2 to 3 days you may need to take a laxative (medication to help you poo) to help you have regular bowel movements. Ask your health care team what to do.</li> </ul> <p>Ask your health care team for the <a href="#">Constipation</a> Pamphlet for more information.</p>	
<p><b>Dizziness</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• You may feel light-headed and like you might faint (pass out).</li> </ul> <p><b>What to do?</b></p> <ul style="list-style-type: none"> <li>• Lay down right away so you do not fall.</li> <li>• Slowly get up and start moving once you feel better.</li> <li>• Do not drive a motor vehicle or use machinery if you feel dizzy.</li> </ul>	Talk to your health care team if it does not improve or if it is severe
<p><b>Neuropathy (Tingling, numb toes or fingers)</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Numbness or tingling of your fingers and toes may happen after starting your treatment.</li> <li>• Sometimes it can be painful and feel like burning sensation, which may be severe.</li> </ul> <p><b>What to do?</b></p> <ul style="list-style-type: none"> <li>• <b>Talk to your health care team if you have symptoms of neuropathy.</b></li> <li>• Numbness and tingling may slowly get better after your treatment ends.</li> </ul> <p>In rare cases, it may continue long after treatment ends. If you continue to have</p>	Talk to your health care team, especially if you have trouble doing tasks like doing up buttons, writing, moving, or if you have severe pain or numbness

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
bothersome symptoms, talk to your health care team for advice.	
<p><b>Trouble Sleeping</b></p> <p>Your medications may cause trouble sleeping. It may get better once your body gets used to the medication or when your treatment ends.</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>You may find it hard to fall asleep or stay asleep.</li> <li>How well you sleep may change over your treatment. For example, you may have several nights of poor sleep followed by a night of better sleep.</li> <li>You may wake up too early or not feel well-rested after a night's sleep.</li> <li>You may feel tired or sleepy during the day.</li> </ul> <p><b>What to do?</b></p> <p>Talk to your health care team if it does not improve or if it is severe</p>	<p>Talk to your health care team if it does not improve or if it is severe</p>
<p><b>Appetite Changes / Weight changes</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>Loss of interest in food or not feeling hungry, or an unexplained increase in appetite</li> <li>Weight loss or weight gain even though you have not changed the way you eat or exercise</li> </ul> <p><b>What to do?</b></p> <ul style="list-style-type: none"> <li>Try to maintain a balanced diet.</li> <li>Ask your health care team to see a dietician to help you control your weight.</li> </ul>	<p>Contact your health care team if no improvement or if severe</p>
<p><b>Too much or too little salt in your body</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>Muscle spasms, cramping, weakness, twitching, or convulsions.</li> <li>Irregular heartbeat, confusion or blood pressure changes.</li> </ul> <p><b>What to do?</b></p>	<p>Get emergency medical help right away for severe symptoms</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<b>Get emergency medical help right away for severe symptoms.</b>	
<p><b>Increased sweating</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Sweating much more than usual in your hands, feet, underarms or face.</li> <li>• Sweating that happens in unusual situations, such as in cooler weather, or without any reason at all.</li> </ul> <p><b>What to do?</b></p> <p><b>To help prevent sweating :</b></p> <ul style="list-style-type: none"> <li>• Avoid triggers such as alcohol, caffeine (tea, coffee, cola), chocolate, hot &amp; spicy food, stress and heat.</li> <li>• Keep cool and dress lightly or in layers that you can remove. Wear clothes made with natural, breathable materials.</li> <li>• Bathe daily.</li> <li>• Drink plenty of liquids unless you are on liquid restriction.</li> <li>• It may improve over time.</li> </ul>	<p>Contact your health care team if no improvement or if severe</p>
<p><b>Mouth sores</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Round, painful, white or gray sores inside your mouth that can occur on the tongue, lips, gums, or inside your cheeks.</li> <li>• In more severe cases they may make it hard to swallow, eat or brush your teeth.</li> <li>• They may last for 3 days or longer.</li> </ul> <p><b>What to do?</b></p> <p><b>To help prevent mouth sores:</b></p> <ul style="list-style-type: none"> <li>• Take care of your mouth by gently brushing and flossing regularly.</li> <li>• Rinse your mouth often with a homemade mouthwash.</li> <li>• To make a homemade mouthwash, mix 1 teaspoonful of baking soda and 1 teaspoonful of salt in 4 cups (1L) of water.</li> <li>• Do not use store-bought mouthwashes, especially those with alcohol, because they may irritate your mouth.</li> </ul> <p><b>If you have mouth sores:</b></p>	<p>Talk to your health care team as soon as you notice mouth or lip sores or if it hurts to eat, drink or swallow</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<ul style="list-style-type: none"> <li>• Avoid hot, spicy, acidic, hard or crunchy foods.</li> <li>• Your doctor may prescribe a special mouthwash to relieve mouth sores and prevent infection.</li> <li>• Talk to your health care team as soon as you notice mouth or lip sores or if it hurts to eat, drink or swallow.</li> </ul> <p>Ask your health care team for the <a href="#">Oral Care (Mouth Care)</a> pamphlet for more information.</p>	

**Other rare, but serious side effects are possible.**

**If you experience ANY of the following, speak to your cancer health care provider or get emergency medical help right away:**

- irregular heartbeat, fainting, chest pain or shortness of breath which worsens with activity
- swelling in your legs, ankles, feet or belly
- limb pain or swelling, hardening of a vein in your limb, sudden loss of the use of your limb(s)
- sudden loss of vision or speech, confusion
- seizures
- signs of an allergic reaction: severe rash, itchiness, swollen face, lip or tongue, chest or throat tightness; may occur during or shortly after the drug is given
- passing little or no urine (pee), foamy urine, or dark urine
- unusual weight gain or unexplained appetite or weight loss
- lower back pain, pain in centre of your belly that may extend to the back
- severe muscle pain or weakness
- yellow skin and/or eyes
- severe tiredness or shortness of breath during mild exertion
- tiny red or purple dots under the skin

## Who do I contact if I have questions or need help?

The most updated information sheet version can be found on <https://www.cancercareontario.ca/drugs>  
 Additional symptom management information is available from <https://www.cancercareontario.ca/symptoms>  
 Created by the CCO Drug Formulary team, with input from the CCO Patient Education team and Patient & Family Advisors.  
 May 2022

My cancer health care provider is: \_\_\_\_\_

During the day I should contact: \_\_\_\_\_

Evenings, weekends and holidays:

**Other Notes:**

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.**May 2022** Revised Medication Information Sheet

For more links on how to manage your symptoms go to [www.cancercareontario.ca/symptoms](http://www.cancercareontario.ca/symptoms).

*The information set out in the medication information sheets, regimen information sheets, and symptom management information (for patients) contained in the Drug Formulary (the "Formulary") is intended to be used by health professionals and patients for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or side effects of a certain drug, nor should it be used to indicate that use of a particular drug is safe, appropriate or effective for a given condition.*



*A patient should always consult a healthcare provider if he/she has any questions regarding the information set out in the Formulary. The information in the Formulary is not intended to act as or replace medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.*