

FINAL DISCHARGE RECOMMENDATIONS

COLPOSCOPY SERVICES

Colposcopist name:

Contact information:

Date:

Patient identifier:

This patient is now discharged from colposcopy. She requires Pap screening by a primary care provider:

- Every three years (routine cervical screening)
- Every year (surveillance)

Re-referral to colposcopy in the future should be guided by her screening results.

According to the Ontario Cervical Screening Program's recommendations, whether or not a woman has been treated, further colposcopic examinations are not required and she can be discharged to primary care if:

HPV testing was not done	HPV testing was done
<input type="checkbox"/> Colposcopy negative AND negative cytology on 3 consecutive visits. Pap screening every 3 years by a primary care provider. <i>These patients are at very low risk for high-grade dysplasia or cervical cancer.</i>	<input type="checkbox"/> HPV test is negative AND normal or low-grade cytology. Pap screening every 3 years by a primary care provider. <i>These patients are at very low risk for high-grade dysplasia or cervical cancer.</i>
<input type="checkbox"/> Colposcopy negative AND any combination of normal or low-grade cytology on 3 consecutive visits. Pap screening every year by a primary care provider. <i>These patients are at slightly elevated risk for high-grade dysplasia or cervical cancer and should be screened annually.</i>	<input type="checkbox"/> HPV test is positive AND normal or low-grade cytology. Pap screening every year by a primary care provider. <i>These patients are at slightly elevated risk for high-grade dysplasia or cervical cancer and should be screened annually.</i>

For further information on screening and colposcopy recommendations for Ontario see cancercare.on.ca/pcs/screening/cervscreening/hcpresources.

, MD, Colposcopist