

Medication Information Sheet

everolimus (E-ver-OH-li-mus)

This document provides general information about your medication. It does not replace the advice of your health care professional. Always discuss your therapy with your health care professional and refer to the package insert for more details.

Other Name: Afinitor®

Appearance: tablet In various strengths

What is this medication for?

For treating certain types of kidney and breast cancer, or other cancers which come from the pancreas, lung or stomach (neuroendocrine tumours).

What should I do before I have this medication?

- Tell your health care team if you have or had significant medical condition(s) such as:
 - ◊ lung, kidney, or liver problems (including hepatitis),
 - ◊ diabetes,
 - ◊ high cholesterol,
 - ◊ active infection(s),
 - ◊ upcoming surgery,
 - ◊ or have any allergies.
- This drug contains a small amount of lactose. If you cannot tolerate lactose, talk to your doctor.
- People with cancer have a higher risk of getting other cancers or developing blood clots. Some cancer medications may increase these risks, especially if used for a long period of time. Discuss any concerns about this medication with your health care team.

How will this medication affect sex, pregnancy and breastfeeding?

- The use of this medication in men or women may cause harm to the unborn baby if pregnancy occurs. Let your health care team know if you or your partner is pregnant, becomes pregnant during treatment, or if you are breastfeeding
- If there is ANY chance that you or your partner may become pregnant, you and your partner together must: ► **Use 2 effective forms of birth control at the same time** while taking this drug: Keep using birth control until **8 weeks** after the last dose. Discuss with your healthcare team.

*The most updated version and more symptom control information can be found on: <http://www.cancercare.on.ca/druginfo>

Prepared with input from the Cancer Care Ontario-Medication Information Sheets Working Group.

December 2017

- Do not breastfeed while taking this drug and for **2 weeks** after the last dose.
- This medication may affect fertility (ability to get pregnant)

How is this medication given?

- Take it exactly as directed by your doctor. Make sure you understand the instructions.
- Do not crush or chew the tablets.
- Take the dose at about the same time each day, preferably in the morning, either always with food or always without food.
- If you miss a dose, take it if it is within 6 hours of the missed dose, otherwise skip this and take the next dose on the following day at its usual scheduled time. Do not double the dose to make up for the forgotten one.

What else do I need to know while on this medication?

- This medication can interact with other medications and can result in the treatment not working as well or cause severe side effects.
- Make sure your health care team knows about all your medications (prescription, over-the-counter, herbals and supplements). Check with your health care team before starting or stopping any of them.
- Do not eat or drink grapefruit, starfruit, Seville oranges or their juices (or products that contain these) while on this treatment. They may increase side effects.
- For mild aches and pain:
 - You may take acetaminophen (Tylenol®) tablets. Ask your health care team about the right dose for you.
 - Ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or aspirin (acetylsalicylic acid, ASA), including low dose aspirin for heart conditions, may increase your chance of bleeding.
 - Talk to your health care team before you start or stop these medications.
 - If you feel unwell, take your temperature before taking any of these medications. They may hide a fever.
 - **Talk to your health care team or go to the closest emergency room right away if you have a fever.** See the [Fever](#) pamphlet for more information.
- Drinking alcohol and smoking during your treatment may increase some side effects and make your medication less effective. Speak to your health care team about smoking and drinking alcohol while on treatment.

How should I safely store this medication?

- Keep this medication in the original packaging at room temperature in a dry place, away from heat and light. Keep out of sight and reach of children and pets.
- Do not throw out any unused medications at home. Bring them to your pharmacy to be thrown away safely.

What are the side effects of this medication?

You may not have all of the side effects below. You may have side effects that are not listed.

Side effects and what to do	When to contact doctor?
Common Side Effects (25 to 49 out of 100 people)	
<p>Mouth sores</p> <p>You may have round, painful, white or gray sores inside your mouth. They can occur on the tongue, lips, gums, or inside your cheeks. In more severe cases they may make it hard to swallow, eat or to brush your teeth. They usually last 1 to 2 weeks.</p> <p>To help prevent mouth sores:</p> <ul style="list-style-type: none"> • Take care of your mouth by gently brushing and flossing regularly. • Rinse your mouth often. Do not use mouthwashes with alcohol. • Instead, try a homemade mouthwash: • Mix 1 teaspoonful of baking soda and 1 teaspoonful of salt in 4 cups (1L) of water. <p>If you have mouth sores:</p> <ul style="list-style-type: none"> • Check with your health care team as soon as you notice mouth or lip sores or if it hurts to swallow. • Avoid hot, spicy, acidic, hard or crunchy foods. Your doctor may prescribe a mouthwash to relieve mouth sores and prevent infection. <p>See the Mouth Care pamphlet for more information.</p>	<p>Contact your health care team as soon as possible</p>
<p>Fatigue (tiredness)</p> <ul style="list-style-type: none"> • Be active and aim to get 30 minutes of moderate exercise (you are able to talk comfortably while exercising) on most days. 	<p>Contact your health care team if no improvement or if severe</p>

*The most updated version and more symptom control information can be found on: <http://www.cancercare.on.ca/druginfo>
 Prepared with input from the Cancer Care Ontario-Medication Information Sheets Working Group.
 December 2017

Side effects and what to do	When to contact doctor?
Common Side Effects (25 to 49 out of 100 people)	
<ul style="list-style-type: none"> • Check with your health care team before starting any new exercise. • Pace yourself, do not rush. Put off less important activities. Rest when you need to. • Eat well and stay hydrated by drinking at least 6 to 8 glasses of water or other liquids every day (unless your doctor told you to drink more or less). • Avoid driving or using machinery if you are feeling tired <p>See our Fatigue pamphlet for more information.</p>	
<p>Cough; feeling short of breath</p> <p>You may have cough and feel short of breath without any signs of infection, such as a sore throat or a stuffed nose.</p>	<p>Contact your health care team if no improvement or if severe</p>
<p>Diarrhea</p> <p>May happen days to weeks after you get your treatment.</p> <p>If you have diarrhea :</p> <ul style="list-style-type: none"> • Take anti-diarrhea medication if your health care team prescribed it. • Avoid foods or drinks with artificial sweetener (e.g. chewing gum, “diet” drinks), coffee and alcohol. • Eat many small meals and snacks instead of 2 or 3 large meals. • Drink at least 6 to 8 cups of liquids each day. Talk to your health care team if you can’t drink 6-8 cups of liquids each day when you have diarrhea. You may need special liquids with salt and sugar, called Oral Rehydration Therapy. <p>See the Diarrhea pamphlet for more information.</p>	<p>Contact your health care team if no improvement or if severe</p>
<p>Rash; dry, itchy skin</p> <p>Rash may be severe in some rare cases and cause your skin to blister or peel. If this happens, get emergency medical help right away.</p> <p>To prevent and treat dry skin,</p> <ul style="list-style-type: none"> • Use skin moisturizer. • Protect your skin from the sun and the cold. • Use sunscreen with UVA and UVB protection and a SPF of at least 30. 	<p>Contact your health care team if no improvement or if severe</p>
<p>Nausea and vomiting (generally mild)</p> <p>May occur in hours to days after your treatment.</p>	<p>Contact your healthcare team if no improvement or</p>

*The most updated version and more symptom control information can be found on: <http://www.cancercare.on.ca/druginfo>
 Prepared with input from the Cancer Care Ontario-Medication Information Sheets Working Group.
 December 2017

Side effects and what to do	When to contact doctor?
Common Side Effects (25 to 49 out of 100 people)	
<p>If you have nausea or vomiting:</p> <ul style="list-style-type: none"> • Take anti-nausea medication(s) as prescribed to you by your doctor. • Drink clear liquids and have small meals. Get fresh air and rest. • Do not eat spicy, fried foods or foods with a strong smell. • Limit caffeine (e.g. coffee, tea) and alcohol. • Contact your health care team if the prescribed anti-nausea medications are not helping to control your nausea and vomiting. <p>Also see Nausea & Vomiting pamphlet for more information.</p>	if severe
<p>Abnormal liver lab tests (may be severe)</p> <ul style="list-style-type: none"> • You may have yellowish skin or eyes, unusually dark pee or pain on the right side of your belly. Talk to your health care team if this happens. • Your doctor may monitor your liver regularly with a blood test. 	Contact your health care team if no improvement or if severe
<p>Low appetite</p> <ul style="list-style-type: none"> • You may not feel like eating or you may lose weight. • Try to eat foods that you like and eat small meals throughout the day. • You may need to take meal supplements to help keep your weight up. • Talk to your health care team if you have a low appetite. • See our Loss of appetite pamphlet for more information. 	Contact your health care team if no improvement or if severe
<p>Mild swelling in arms and legs; puffiness</p> <p>To help prevent swelling :</p> <ul style="list-style-type: none"> • Eat a low-salt diet. • Avoid tight fitting clothing. <p>If you have swelling in your legs, keep your feet up when sitting.</p>	Contact your health care team if no improvement or if severe

Side effects and what to do	When to contact doctor?
Less Common Side Effects (10 to 24 out of 100 people)	
<p>Headache; mild joint, muscle pain or cramps</p> <ul style="list-style-type: none"> • Take your pain medication as prescribed by your doctor. • You can take acetaminophen (Tylenol®) tablets as needed for mild aches and pains. Ask your doctor or pharmacist about the right dose for you. • Talk to your doctor or pharmacist first before taking ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or aspirin. These medication may increase bleeding risk. • Rest often and try light exercise as it may help. 	<p>Contact your health care team if no improvement or if severe</p>
<p>High blood sugar</p> <ul style="list-style-type: none"> • You may feel thirsty and pee more often. • Your doctor may check your blood sugar level. You may be advised to change your diet or take medication to treat high blood sugar. • Check your blood sugar regularly if you have diabetes. 	<p>Contact your health care team as soon as possible</p>
<p>Lung problems</p> <p>You may have a cough, breathing problems, chest pain or cough up blood.</p>	<p>Get emergency medical help right away</p>
<p>Taste changes</p> <ul style="list-style-type: none"> • Eat food that needs less chewing. • Taste foods at different temperatures, since the flavor may change. • Try different forms of foods, like fresh, frozen or canned. • Experiment with non-spicy foods, spices and seasonings. 	<p>Contact your health care team if no improvement or if severe</p>
<p>Unusual bleeding or bruising</p> <p>You may have black stools, cough up blood, blood in your urine, purple or red dots on your skin or bleeding that will not stop.</p> <p>Fever, chills, infection</p> <p>You have a fever if your temperature taken in your mouth (oral temperature) is:</p> <ul style="list-style-type: none"> • 38.3°C (100.9°F) or higher at any time OR • 38.0°C (or 100.4°F) or higher for at least one hour. <p>While you are getting treatment:</p> <ul style="list-style-type: none"> • Keep a digital thermometer at home and take your temperature if you feel hot or unwell (for example, chills). • Avoid taking medications that treat a fever before you take your temperature (for example, Tylenol®, acetaminophen, Advil® or ibuprofen) 	<p>Get emergency medical help right away</p>

*The most updated version and more symptom control information can be found on: <http://www.cancercare.on.ca/druginfo>
 Prepared with input from the Cancer Care Ontario-Medication Information Sheets Working Group.
 December 2017

Side effects and what to do	When to contact doctor?
Less Common Side Effects (10 to 24 out of 100 people)	
<p>as they may hide a fever.</p> <ul style="list-style-type: none"> • Do not eat or drink anything hot or cold right before taking your temperature. • Wash your hands often. • Check with your doctor before getting any vaccines, surgeries, medical procedures or visiting your dentist. <p>If you have a fever, talk to your health care team or go to the closest emergency room. See our Neutropenia (Low white blood cell count) pamphlet for more information.</p>	

Other rare, but serious side effects are possible.

If you experience ANY of the following, speak to your cancer health care provider or get emergency medical help right away:

- signs of an allergy such as fever, itchiness, rash, swollen lips, face or tongue, chest and throat tightness.
- pain, swelling and hardening of the vein in your arm or leg
- irregular heartbeat, shortness of breath, chest pain, fainting spells or swelling in your legs, ankles and belly
- severe belly pain, bloating or feeling of fullness and severe constipation
- severe muscle pain or weakness and dark-coloured pee
- lower back pain, swelling, pee less than usual and have unusual weight gain
- sudden and severe tiredness and shortness of breath during mild activity
- wounds that do not heal well

For more links on how to manage your symptoms go to www.cancercare.on.ca/symptoms.

The information set out in the medication information sheets, regimen information sheets, and symptom management information (for patients) contained in the Drug Formulary (the "Formulary") is intended to be used by health professionals and patients for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or side effects of a certain drug, nor should it be used to indicate that use of a particular drug is safe, appropriate or effective for a given condition.

A patient should always consult a healthcare provider if he/she has any questions regarding the information set out in the Formulary. The information in the Formulary is not intended to act as or replace medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.

*The most updated version and more symptom control information can be found on: <http://www.cancercare.on.ca/druginfo>
Prepared with input from the Cancer Care Ontario-Medication Information Sheets Working Group.
December 2017

