

Regimen Monograph

[Regimen Name](#) | [Drug Regimen](#) | [Cycle Frequency](#) | [Premedication and Supportive Measures](#) | [Administrative Information](#) | [References](#) | [Other Notes](#) | [Disclaimer](#)

A - Regimen Name

MFOLFOX6 Regimen

Folinic Acid (Leucovorin)-Fluorouracil-Oxaliplatin

Disease Site Gastrointestinal
 Esophagus
 Gastric / Stomach

Intent Adjuvant
 Palliative

Regimen Category **Evidence-informed :**

Regimen is considered appropriate as part of the standard care of patients; meaningfully improves outcomes (survival, quality of life), tolerability or costs compared to alternatives (recommended by the Disease Site Team and national consensus body e.g. pan-Canadian Oncology Drug Review, pCODR). Recommendation is based on an appropriately conducted phase III clinical trial relevant to the Canadian context OR (where phase III trials are not feasible) an appropriately sized phase II trial. Regimens where one or more drugs are not approved by Health Canada for any indication will be identified under Rationale and Use.

This **Regimen Abstract** is an **abbreviated** version of a Regimen Monograph and contains only top level information on usage, dosing, schedule, cycle length and special notes (if available). Information in regimen abstracts is accurate to the extent of the ST-QBP regimen master listings, and has not undergone the full review process of a regimen monograph. Full regimen monographs will be published for each ST-QBP regimen as they are developed.

[back to top](#)

B - Drug Regimen

oxaliplatin	85 mg /m ²	IV in 500mL D5W over 120 minutes	Day 1
leucovorin	400 mg /m ²	IV diluted in D5W over 120 minutes (concurrently with oxaliplatin)	Day 1
fluorouracil THEN	400 mg /m ²	IV bolus, after leucovorin	Day 1
fluorouracil	2400 mg /m ²	IV continuous infusion over 46 hours (single dose)	Start on Day 1

[back to top](#)**C - Cycle Frequency****REPEAT EVERY 14 DAYS**

Adjuvant: For a usual total of 12 cycles unless disease progression or unacceptable toxicity occurs

Palliative: Until disease progression or unacceptable toxicity

[back to top](#)**D - Premedication and Supportive Measures**

Antiemetic Regimen: Moderate

**Febrile Neutropenia
Risk:** Low

Other Supportive Care:

Also refer to [CCO Antiemetic Recommendations](#).

[back to top](#)

J - Administrative Information

Approximate Patient Visit	3 hours
Pharmacy Workload (average time per visit)	38.381 minutes
Nursing Workload (average time per visit)	69.167 minutes

[back to top](#)

K - References

Noh SH, Park SR, Yang HK, et al. Adjuvant capecitabine plus oxaliplatin for gastric cancer after D2 gastrectomy (CLASSIC): 5-year follow-up of an open-label, randomised phase 3 trial. *Lancet Oncol* 2014;15(12):1389-96.

Al-Batran SE, Hartmann JT, Probst S, et al. Phase III trial in metastatic gastroesophageal adenocarcinoma with fluorouracil, leucovorin plus either oxaliplatin or cisplatin: a study of the Arbeitsgemeinschaft Internistische Onkologie. *J Clin Oncol* 2008;26(9):1435-42.

PEBC Advice Documents or Guidelines

- [Systemic Therapy for Advanced Gastric and Gastro-Esophageal Carcinoma](#)

April 2023 Updated DPD deficiency and fluorouracil antidote information in the Other Notes section

[back to top](#)

L - Other Notes

DPD Deficiency Testing and Guidance

Patients should be tested for DPD deficiency before starting treatment with fluorouracil. Refer to the [DPD Deficiency Guidance for Clinicians](#) for more information.

In patients with unrecognized DPD deficiency, acute, life-threatening toxicity may occur; if acute grade 2-4 toxicity develops, treatment should be stopped immediately and permanent discontinuation considered based on clinical assessment of the toxicities.

Antidote for Fluorouracil Overdose:

Uridine triacetate is a prodrug of uridine and is a specific antidote for treating fluorouracil

overdose or severe early onset toxicities. If available, consider administering as soon as possible (i.e. within 96 hours) for suspected overdose. If not available, treatment is symptomatic and supportive.

For usage approval and supply, contact Health Canada's [Special Access Program](#) (SAP) (Phone: 613-941-2108. On-call service is available for emergencies). Uridine triacetate (Vistogard®) is supplied by its manufacturer in the United States (Wellstat Therapeutics).

The recommended dosing and administration for **uridine triacetate** in patients ≥ 18 years is:

- 10 grams (1 packet of coated granules) orally every 6 hours for 20 doses in total, without regards to meals.
- Granules should not be chewed. They should be mixed with 3 to 4 ounces of soft foods such as applesauce, pudding or yogurt.
- The dose should be ingested within 30 minutes of preparation, followed by at least 4 ounces of water.
- Refer to the prescribing information on dose preparation for NG-tube or G-tube use.

Additional resources on the management of fluorouracil infusion overdose:

- [Management of Fluorouracil Infusion Overdose Guideline](#) (Alberta Health Services)
- [Management of Fluorouracil Infusion Overdose at the BCCA - Interim Guidance](#) (BC Cancer Agency)

[back to top](#)

M - Disclaimer

Regimen Abstracts

A Regimen Abstract is an abbreviated version of a Regimen Monograph and contains only top level information on usage, dosing, schedule, cycle length and special notes (if available). It is intended for healthcare providers and is to be used for informational purposes only. It is not intended to constitute or be a substitute for medical advice, and all uses of the Regimen Abstract are subject to clinical judgment. Such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability, and Cancer Care Ontario disclaims all liability for the use of this information, and for any claims, actions, demands or suits that arise from such use.

Information in regimen abstracts is accurate to the extent of the ST-QBP regimen master listings, and has not undergone the full review process of a regimen monograph. Full regimen monographs will be published for each ST-QBP regimen as they are developed.

Regimen Monographs

Refer to the [New Drug Funding Program](#) or [Ontario Public Drug Programs](#) websites for the most up-to-date public

funding information.

The information set out in the drug monographs, regimen monographs, appendices and symptom management information (for health professionals) contained in the Drug Formulary (the "Formulary") is intended for healthcare providers and is to be used for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or adverse effects of a particular drug, nor should it be construed to indicate that use of a particular drug is safe, appropriate or effective for a given condition. The information in the Formulary is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.

The format and content of the drug monographs, regimen monographs, appendices and symptom management information contained in the Formulary will change as they are reviewed and revised on a periodic basis. The date of last revision will be visible on each page of the monograph and regimen. Since standards of usage are constantly evolving, it is advised that the Formulary not be used as the sole source of information. It is strongly recommended that original references or product monograph be consulted prior to using a chemotherapy regimen for the first time.

Some Formulary documents, such as the medication information sheets, regimen information sheets and symptom management information (for patients), are intended for patients. Patients should always consult with their healthcare provider if they have questions regarding any information set out in the Formulary documents.

While care has been taken in the preparation of the information contained in the Formulary, such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability.

CCO and the Formulary's content providers shall have no liability, whether direct, indirect, consequential, contingent, special, or incidental, related to or arising from the information in the Formulary or its use thereof, whether based on breach of contract or tort (including negligence), and even if advised of the possibility thereof. Anyone using the information in the Formulary does so at his or her own risk, and by using such information, agrees to indemnify CCO and its content providers from any and all liability, loss, damages, costs and expenses (including legal fees and expenses) arising from such person's use of the information in the Formulary.

[back to top](#)