

Drug Monograph

[Drug Name](#) | [Mechanism of Action and Pharmacokinetics](#) | [Indications and Status](#) | [Adverse Effects](#) | [Dosing](#) | [Administration Guidelines](#) | [Special Precautions](#) | [Interactions](#) | [Recommended Clinical Monitoring](#) | [Supplementary Public Funding](#) | [References](#) | [Disclaimer](#)

A - Drug Name

PACLitaxel

COMMON TRADE NAME(S): Taxol®

[back to top](#)

B - Mechanism of Action and Pharmacokinetics

Paclitaxel promotes assembly of microtubules and stabilizes them against depolymerization. It also inhibits cell replication by blocking cells in the late G2 and/or M phases of the cell cycle. Paclitaxel is obtained via a semi-synthetic process from *Taxus baccata*.

Distribution

Extensive extravascular distribution and/or tissue binding.

Cross blood brain barrier? no

PPB 89 %

Metabolism

Hepatic metabolism (CYP 2C8 and CYP 3A4) and biliary secretion. Non-linear pharmacokinetics.

Active metabolites no information found

Inactive metabolites Hydroxylated metabolites

Elimination

High concentrations found in bile; 71% excreted in feces in 120 hours (5% unchanged)

Urine 1.3 to 12.7 % as unchanged drug.

Half-life

9.9 hours (3 hr infusion)

[back to top](#)

C - Indications and Status

Note: There are multiple generic brands, and not all of the indications listed below are contained in each product monograph. However, the indication is listed if approved for at least one generic brand

Health Canada Approvals:

- Breast cancer (adjuvant and second-line metastatic)
- Non-small cell lung cancer (first-line advanced)
- Ovarian cancer (first-line combination, second-line metastatic)
- AIDS-related Kaposi's sarcoma (advanced; refractory to liposomal anthracycline)

Other Uses:

- Breast cancer (neoadjuvant)
- Gastrointestinal cancer (anal, gastroesophageal)
- Genitourinary cancer (penile, prostate, bladder, testicular)
- Gynecological cancer (cervical, germ cell, gestational trophoblastic disease, vulvar, gynecological sarcoma, endometrial, fallopian tube, primary peritoneal)
- Head and neck cancer
- Thyroid cancer
- Thymoma
- Ewing's sarcoma
- Skin cancer (Merkel cell, Melanoma)
- Cancer of unknown primary origin

[back to top](#)

D - Adverse Effects

Emetogenic Potential: Low

Extravasation Potential: Irritant

The following data were based on clinical trials in ovarian and breast cancer patients, who were

treated with single agent paclitaxel 175 mg/m² IV over 3 hours.

ORGAN SITE	SIDE EFFECT* (%)	ONSET**
Cardiovascular	Arrhythmia (3%) (transient, including bradycardia)	I
	Arterial thromboembolism (rare)	E
	Cardiotoxicity (rare)	E D
	ECG changes (14%)	E
	Hypotension (11%)	I
	Venous thromboembolism (rare)	E
Dermatological	Alopecia (93%) (rarely permanent)	E
	Nail disorder (2%)	E D
	Radiation recall reaction (rare)	I
	Rash (rare- may be severe)	I E
Gastrointestinal	Dehydration	E
	Diarrhea (25%)	I E
	GI obstruction (rare)	E
	GI perforation (rare)	E
	Mucositis (20%)	E
	Nausea, vomiting (44%)	I
	Typhlitis (rare)	I
General	Edema (21%)	E
	Fatigue (17%)	E
Hematological	Myelosuppression ± infection, bleeding (grade 4 neutropenia 27%)	E
Hepatobiliary	↑ LFTs (18%) (severe- rare)	E
	Pancreatitis (rare)	E
Hypersensitivity	Hypersensitivity (40%) (severe 1%)	I
Injection site	Injection site reaction (4%) (may be severe)	I
Musculoskeletal	Musculoskeletal pain (54%) (severe 12%)	E
Neoplastic	Secondary malignancy (rare)	L
Nervous System	Ataxia (rare)	E
	Autonomic neuropathy (rare)	E
	Encephalopathy (rare)	E
	Optic neuritis / Ototoxicity (rare)	E

	Peripheral neuropathy (64%) (severe 4%)	E
	Seizure (rare)	E
	Syncope (rare)	E
Ophthalmic	Cystoid macular edema (rare)	E
	Eye disorders (visual disturbances- rare)	E
	Optic nerve disorder (rare)	E
Respiratory	Pneumonitis (rare)	E

* "Incidence" may refer to an absolute value or the higher value from a reported range.
 "Rare" may refer to events with < 1% incidence, reported in post-marketing, phase 1 studies, isolated data or anecdotal reports.

** I = *immediate* (onset in hours to days) E = *early* (days to weeks)
 D = *delayed* (weeks to months) L = *late* (months to years)

The most common adverse events are alopecia, peripheral neuropathy, musculoskeletal pain, nausea/vomiting, hypersensitivity, myelosuppression, diarrhea, edema, mucositis, ↑ LFTs.

Myelosuppression is dose and schedule-dependent but is not cumulative, with neutropenia being less common when paclitaxel was given as a 3-hour infusion as compared to a 24-hour infusion schedule. Toxicity may be more severe in HIV patients, especially infection (febrile neutropenia and opportunistic infections) and neutropenia.

Congestive heart failure (including LVEF decrease) has been reported in patients who have received other chemotherapy agents, especially anthracyclines.

Hypersensitivity reactions typically occur in early treatment courses and within the first hour of infusion. Dyspnea, flushing, chest pain and tachycardia were the most frequent manifestations. Reactions are neither dose-related nor dependent on prior exposure to paclitaxel, and may be caused by histamine release mediated by the Cremophor EL diluent. Because of the significant risk of hypersensitivity reactions, the patient must be monitored closely; a physician must be readily available, as well as emergency medications and resuscitation equipment. Anaphylaxis and severe hypersensitivity reactions (hypotension, angioedema, generalized urticaria) occur in 2% of patients and may rarely be fatal.

Myalgia and/or arthralgia tend to appear 2-3 days after paclitaxel administration and resolve within a few days, and do not appear to be dose-related. Non-steroidal anti-inflammatory drugs are successful in relieving these symptoms.

Peripheral neuropathy may be dose-limiting and is dose-related and cumulative. Common symptoms include numbness, tingling and/or burning pain in a glove-and-stocking distribution. Mild symptoms usually improve or resolve completely within several months after discontinuation of therapy. Pre-existing neuropathies are not a contraindication to treatment with paclitaxel.

Central neurotoxicity may occur and may be severe, especially in children treated at high dosage.

Hypotension and bradycardia have been observed during paclitaxel infusion and are usually

asymptomatic, but not dose or schedule-dependent. Severe cardiovascular events, including death, have been reported; if these occur, appropriate action should be taken and the dose interrupted/discontinued; continuous electrocardiographic monitoring should be performed if patient receives subsequent paclitaxel therapy.

Cystoid macular edema (CME) has been reported in paclitaxel-treated patients, as well as with other taxanes. Patients who present with impaired vision during treatment should undergo a prompt ophthalmologic examination. Taxane-associated CME may not be associated with vascular leakage and is usually reversible upon taxane discontinuation. Treatment for CME may be required in some cases.

Paclitaxel has the potential to enhance radiation injury to tissues. While often called **radiation recall reactions**, the timing of the radiation may be before, concurrent with or even after the administration of paclitaxel. Recurrent injury to a previously radiated site may occur weeks to months following radiation. Recall skin reactions at a site of previous extravasation, after paclitaxel administration at a different site, have been reported rarely.

[back to top](#)

E - Dosing

Refer also to protocol by which patient is being treated.

Pre-medications* (prophylaxis for infusion reaction):

Pre-Medications for Q3W paclitaxel:

- Dexamethasone 20 mg PO 12-and 6-hours OR Dexamethasone 20 mg IV 30 minutes pre-infusion[†]
- Diphenhydramine 25-50 mg IV/PO 30-60 minutes pre-infusion
- Ranitidine 50 mg IV OR Famotidine 20 mg IV 30-60 minutes pre-infusion

Pre-Medications for weekly paclitaxel: (To be given 30-60 minutes prior to paclitaxel infusion)

- Dexamethasone 10 mg IV, starting in cycle 1
- Diphenhydramine 25-50 mg IV/PO
- Ranitidine 50 mg IV OR Famotidine 20 mg IV

* Consider **discontinuing** pre-medications for paclitaxel if there was no IR in the first 2 doses.

[†] Oral and IV dexamethasone are both effective at reducing overall IR rates. Some evidence suggests that oral dexamethasone may be more effective for reducing severe reactions; however, adverse effects and compliance remain a concern.

Adults:

Start treatment only in patients with neutrophils $\geq 1.5 \times 10^9/L$ ($\geq 1 \times 10^9/L$ in AIDS-related Kaposi's sarcoma) and platelet counts $\geq 100 \times 10^9/L$

Intravenous: 175 mg/m² over 3 hours every 3 weeks

AIDS-related Kaposi's Sarcoma:

Intravenous: 135 mg/m² over 3 hours every 3 weeks
OR

AIDS-related Kaposi's Sarcoma:

Intravenous: 100 mg/m² over 3 hours every 2 weeks

Dosage with Toxicity:

Worst toxicity in previous cycle	Dose of paclitaxel
Febrile neutropenia Grade 4 ANC ≥ 5 -7 days Grade 4 thrombocytopenia	↓ by 20%*
Grade 3 neurotoxicity or other toxicity	↓ by 20%*
Grade 4 neurotoxicity or other toxicity, any grade cystoid macular edema	Discontinue
*Patients should not be retreated with paclitaxel until neutrophils $\geq 1.5 \times 10^9/L$ ($\geq 1.0 \times 10^9/L$ in AIDS-related Kaposi's sarcoma) and platelet counts $\geq 100 \times 10^9/L$ and other toxicity has recovered to \leq grade 2	

Management of Infusion-related reactions:

Also refer to the CCO guideline for detailed description of [Management of Cancer Medication-Related Infusion Reactions](#).

Grade	Management	Re-challenge
1 or 2	<ul style="list-style-type: none"> Stop or slow the infusion rate. Manage the symptoms. <p>Restart:</p> <ul style="list-style-type: none"> After symptom resolution, restart with pre-mediations ± reduced infusion rate. 	<ul style="list-style-type: none"> Consider re-challenge with pre-mediations and at a reduced infusion rate. After 2 subsequent IRs, consider replacing with a different taxane. Give intensified pre-mediations and reduce the infusion rate. May consider adding oral montelukast ± oral acetylsalicylic acid.
3 or 4	<ul style="list-style-type: none"> Stop treatment. Aggressively manage symptoms. 	<ul style="list-style-type: none"> Re-challenge is discouraged, especially if vital signs have been affected. Consider desensitization if therapy is necessary. There is insufficient evidence to recommend substitution with another taxane at re-challenge. High cross-reactivity rates have been reported.

Dosage with Hepatic Impairment:

Caution and dose reduction advised in patients with moderate to severe hepatic impairment.

Patients with hepatic impairment may be at risk of toxicity, especially severe myelosuppression.

Suggested are:

Bilirubin		AST/ALT	Dose (% usual dose)
≤1.25 x ULN	And	2-10 x ULN	75%
1.26 to 2.5 x ULN	And	<10x ULN	40%
2.6 to 4 x ULN	And	<10x ULN	25%
>4 x ULN	And/Or	≥10 x ULN	Consider risk-benefit or Omit

Dosage with Renal Impairment:

No adjustment required, but may consider for patients with HIV-AIDS if creatinine $\geq 2 \times$ ULN

Dosage in the elderly:

No adjustment required, but elderly patients are more at risk for severe toxicity.

Children:

Safety and efficacy have not been established. Children may be at a higher risk of severe and sometimes fatal neurologic toxicity, especially with high doses, possibly related to the ethanol content of paclitaxel infusions.

[back to top](#)

F - Administration Guidelines

- In order to minimize patients' exposure to DEHP leaching from PVC bags or sets, use polyolefin or polypropylene infusion bags and polyethylene-lined administration sets (with a 0.22 micron in-line filter).
- Dilute in 500-1000 mL Normal Saline or 5% Dextrose, in a final concentration of 0.3-1.2 mg/mL and infuse over 3 hours.
- For weekly dosing, may be infused over 1 hour - mix in 250mL bag as above (not approved by manufacturer).
- Extended infusion of paclitaxel is not recommended as primary prophylaxis to reduce paclitaxel IRs.
- Paclitaxel should be given before cisplatin, if given in combination (refer to drug interactions section)
- Excessive shaking, agitation, or vibration may induce precipitation and should be avoided
- Precipitation may rarely occur with infusions longer than 3 hours.

Also refer to the CCO guideline for detailed description of [Management of Cancer Medication-Related Infusion Reactions](#).

[back to top](#)

G - Special Precautions

Contraindications:

- Patients with a history of severe hypersensitivity reactions to paclitaxel or other drugs formulated in Cremophor EL (polyethoxylated castor oil)
- Patients with severe baseline neutropenia ($<1.5 \times 10^9/L$; $< 1 \times 10^9/L$ for patients with AIDS-related Kaposi's)

Other Warnings/Precautions:

- Paclitaxel contains ethanol, and is administered with agents such as antihistamines which cause drowsiness. Patients should be cautioned regarding driving and the use of machinery.

Pregnancy and Lactation:

- Embryotoxicity: Yes
- Fetotoxicity: Yes
Paclitaxel is not recommended for use in pregnancy. Adequate contraception should be used by both sexes during treatment, and for at least 6 months after the last dose.
- Excretion into breast milk: Probable
Breastfeeding is not recommended.
- Fertility effects: Yes

[back to top](#)

H - Interactions

The drugs listed in this table are based on drug interaction case reports, pharmacokinetic studies, or potential interactions. Paclitaxel clearance was not affected by cimetidine administration pre-treatment.

AGENT	EFFECT	MECHANISM	MANAGEMENT
Cisplatin	↑ myelosuppression if cisplatin given prior (within hours) to paclitaxel	↓ paclitaxel clearance by 33%	Give paclitaxel before cisplatin
Cisplatin	May ↑ risk of renal failure in gynaecological cancers	Unknown	Caution

Doxorubicin (after prolonged paclitaxel infusions)	↑ neutropenia and stomatitis	Higher plasma levels of doxorubicin and doxorubicinol	Caution
Epirubicin	↑ systemic exposure to epirubicin or its metabolites. May be schedule-dependent.	Taxane or cremophor EL possibly compete with epirubicin for biliary excretion	Caution if used in combination; give epirubicin first
Cyclophosphamide (given after paclitaxel)	↑ myelosuppression	Unknown	Caution
Radiation	Radiation pneumonitis	↑ pulmonary effects	Avoid/caution
Carboplatin (given after paclitaxel)	↓ thrombocytopenia	Unknown	Caution
CYP 2C8 substrates (i.e. paclitaxel, sorafenib, amiodarone)	May ↑/↓ effects of substrates or paclitaxel	Altered metabolism of CYP2C8 substrates or paclitaxel	Caution
CYP 3A4 substrates (i.e., verapamil, etoposide, dexamethasone, vincristine)	May ↑/↓ effects of substrates or paclitaxel	Altered metabolism of CYP3A4 substrates or paclitaxel	Caution
Inducers of CYP 2C8 (i.e., phenobarbital)	May ↓ paclitaxel levels and effects	↑ metabolism of paclitaxel	Caution
CYP 2C8 inhibitors (i.e. gemfibrozil, montelukast)	May ↑ paclitaxel levels and effects	↓ metabolism of paclitaxel	Caution
CYP3A4 inducers (i.e. phenytoin, rifampin, dexamethasone, carbamazepine, phenobarbital, St. John's Wort, etc)	May ↓ paclitaxel levels and effects	↑ metabolism of paclitaxel	Caution
CYP3A4 inhibitors (i.e. ketoconazole, clarithromycin, ritonavir, fruit or juice from	May ↑ paclitaxel levels and effects	↓ metabolism of paclitaxel	Caution

grapefruit, Seville
oranges or
starfruit)

[back to top](#)

I - Recommended Clinical Monitoring

Treating physicians may decide to monitor more or less frequently for individual patients but should always consider recommendations from the product monograph.

Recommended Clinical Monitoring

Monitor Type	Monitor Frequency
CBC	Baseline and before each visit
Liver function tests	Baseline and before each cycle
Renal function tests	Baseline and as clinically indicated
Blood pressure and pulse	Per usual institutional protocol; also during infusion (more frequently during the first hour)
Continuous cardiac monitoring in patients who developed serious conduction abnormalities	During subsequent infusions
Ophthalmology if visual impairment	As clinically indicated
Clinical assessment of bleeding, infection, diarrhea, musculoskeletal, neurologic (sensory), hypersensitivity, respiratory, thromboembolism	At each visit

Grade toxicity using the current [NCI-CTCAE \(Common Terminology Criteria for Adverse Events\) version](#)

Suggested Clinical Monitoring

Monitor Type	Monitor Frequency
Renal function tests (AIDS-related Kaposi's sarcoma)	Baseline and before each dose

[back to top](#)

K - References

Cancer Drug Manual (the Manual), 1994, British Columbia Cancer Agency (BCCA).

Danesi R, Innocenti F, Fogli S, et al. Pharmacokinetics and pharmacodynamics of combination chemotherapy with paclitaxel and epirubicin in breast cancer patients. *Br J Clin Pharmacol* 2002; 53: 508–18.

Clin-Info: e-Compendium of Pharmaceuticals and Specialties. Accessed October 27, 2009.

Esposito M, Venturini M, Vannozzi MO, et al. Comparative effects of paclitaxel and docetaxel on the metabolism and pharmacokinetics of epirubicin in breast cancer patients. *J Clin Oncol* 1999; 17:1132-40.

Floyd J, Mirza I, Sachs B, et al. Hepatotoxicity of chemotherapy. *Semin Oncol* 2006;33(1):50-67.

Fogli S, Danesi R, Gennari A, et al. Gemcitabine, epirubicin and paclitaxel: pharmacokinetic and pharmacodynamic interactions in advanced breast cancer. *Annals of Oncology* 2002; 13: 919–27.

Gill PS, Tulpule, A, Espina BM, et al. Paclitaxel is safe and effective in the treatment of advanced AIDS-Related Kaposi's sarcoma. *J Clin Oncol* 1999; 17: 1876-83.

Grasselli G, Vigano L, Capri G, et al. Clinical and pharmacologic study of the epirubicin and paclitaxel combination in women with metastatic breast cancer. *J Clin Oncol* 2001; 19:2222-31.

King PD and Perry MC. Hepatotoxicity of chemotherapy. *The Oncologist* 2001;6:162-176.

McEvoy GK, editor. AHFS Drug Information 2009. Bethesda: American Society of Health-System Pharmacists, p. 1191-205.

Prevezas C, Matard B, Pinquier L, et al. Irreversible and severe alopecia following docetaxel or paclitaxel cytotoxic therapy for breast cancer. *British Journal of Dermatology* 2009; 160(4): 883-5.

Product Monograph: Taxol® (paclitaxel) Bristol-Myers Squibb Inc (Canada), February 22, 2010.

Product Monograph: Paclitaxel for injection. Hospira Healthcare Corp. (Canada), October 28, 2013.

Superfin D, Iannucci AA, Davies AM, et al. Commentary: Oncologic drugs in patients with organ dysfunction: a summary. *Oncologist* 2007;12(9):1070-83.

Venturini M, Lunardi G, Del Mastro L, et al. Sequence effect of epirubicin and paclitaxel treatment on pharmacokinetics and toxicity. *J Clin Oncol* 2000; 18: 2116-25.

April 2023 removed NDFP forms

[back to top](#)

L - Disclaimer

Refer to the [New Drug Funding Program](#) or [Ontario Public Drug Programs](#) websites for the most up-to-date public funding information.

The information set out in the drug monographs, regimen monographs, appendices and symptom management information (for health professionals) contained in the Drug Formulary (the "Formulary") is intended for healthcare providers and is to be used for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or adverse effects of a particular drug, nor should it be construed to indicate that use of a particular drug is safe, appropriate or effective for a given condition. The information in the Formulary is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.

The format and content of the drug monographs, regimen monographs, appendices and symptom management information contained in the Formulary will change as they are reviewed and revised on a periodic basis. The date of last revision will be visible on each page of the monograph and regimen. Since standards of usage are constantly evolving, it is advised that the Formulary not be used as the sole source of information. It is strongly recommended that original references or product monograph be consulted prior to using a chemotherapy regimen for the first time.

Some Formulary documents, such as the medication information sheets, regimen information sheets and symptom management information (for patients), are intended for patients. Patients should always consult with their healthcare provider if they have questions regarding any information set out in the Formulary documents.

While care has been taken in the preparation of the information contained in the Formulary, such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability.

CCO and the Formulary's content providers shall have no liability, whether direct, indirect, consequential, contingent, special, or incidental, related to or arising from the information in the Formulary or its use thereof, whether based on breach of contract or tort (including negligence), and even if advised of the possibility thereof. Anyone using the information in the Formulary does so at his or her own risk, and by using such information, agrees to indemnify CCO and its content providers from any and all liability, loss, damages, costs and expenses (including legal fees and expenses) arising from such person's use of the information in the Formulary.

[back to top](#)

