

Medication Information Sheet

pegylated liposomal DOXOrubicin (PEG-y-late-ted Lip-o-

SO-mal Docs-oh-RUBE-i-sin)

This document provides general information about your medication. It does not replace the advice of your health care professional. Always discuss your therapy with your health care professional and refer to the package insert for more details.

Other Name: Caelyx®

Appearance: Red coloured liquid mixed into larger bags of fluids

What is this medication for?

- For treating breast cancer, ovarian cancer, and HIV-positive Kaposi's sarcoma.

What should I do before I have this medication?

- Tell your doctor if you have/had significant medical condition(s), especially if you have / had heart disease, or have any allergies.
- Pegylated liposomal doxorubicin may harm the unborn baby.
- Let your doctor know if you are breastfeeding, pregnant or plan to become pregnant.
- People who have cancer or leukemia are at a higher risk of developing other cancers/leukemias (usually some years later) or blood clots. Some cancer medications, such as pegylated liposomal doxorubicin, may increase these risks, especially if used for a prolonged period of time. You should discuss any concerns with your doctor.

How will this medication affect sex, pregnancy and breastfeeding?

- Do not use this drug if you are pregnant. If there is ANY chance that you or your partner may become pregnant, you and your partner together must: ► **Use 2 effective forms of birth control at the same time** while taking this drug. Do not take birth control pills if you have breast cancer. Keep using birth control until **6 months** after the last dose (general recommendation). Discuss with your healthcare team.
- Tell your doctor right away if you or your partner becomes pregnant.
- Do not breastfeed while taking this drug.
- Effects on Fertility: Probable

How is this medication given?

- This drug is given by injection into a vein.

*The most updated version and more symptom control information can be found on: <http://www.cancercare.on.ca/druginfo>

Prepared with input from the Cancer Care Ontario-Medication Information Sheets Working Group.

May 2018

What else do I need to know while on this medication?

- This medication can interact with other medications and can result in the treatment not working as well or cause severe side effects.
- Make sure your health care team knows about all your medications (prescription, over-the-counter, herbals and supplements). Check with your health care team before starting or stopping any of them.
- **For mild aches and pain or fever:**
 - If you feel unwell, take your temperature before taking any medications for pain or fever. They may hide a fever.
 - You may take acetaminophen (Tylenol®) tablets. Ask your health care team about the right dose for you.
 - Ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or aspirin (acetylsalicylic acid, ASA), including low dose aspirin for heart conditions, may increase your chance of bleeding. Talk to your health care team before you start or stop these medications.
 - **Talk to your health care team or go to the closest emergency room right away if you have a fever.** See the [Fever](#) pamphlet for more information.

What are the side effects of this medication?

The following side effects have been seen in people using pegylated liposomal DOXOrubicin with other chemotherapy drugs, so some of these effects may also be related to chemotherapy.

The following side effects are common or severe. You may not have all of the side effects, while others may occur. Discuss with your doctor if you have any unusual or bothersome symptoms.

Side effects and what to do	When to contact doctor?
More Common Side Effects	
<p>Hand-foot syndrome (pain, thinning or reddening palms or feet, including tingling, numbness, peeling)</p> <ul style="list-style-type: none"> • Avoid activities that cause rubbing, pressure or heat exposure to hands and feet (i.e. gripping tools, vigorous washing, hot baths). • Apply moisturizer liberally and often to your hands and feet, especially in the creases. • Wear loose, comfortable footwear and clothes. Rest and try to keep off your feet. 	<p>Contact your health care team if no improvement or if severe</p>

*The most updated version and more symptom control information can be found on: <http://www.cancercare.on.ca/druginfo>
Prepared with input from the Cancer Care Ontario-Medication Information Sheets Working Group.
May 2018

pegylated liposomal DOXOrubicin

Side effects and what to do	When to contact doctor?
More Common Side Effects	
<ul style="list-style-type: none"> • Also see Hand-Foot Syndrome pamphlet.* 	
<p><i>Nausea and vomiting</i></p> <ul style="list-style-type: none"> • Drink clear fluids and avoid large meals. Get fresh air and rest. • Limit spicy, fried foods or foods with a strong smell. • Take anti-nausea drug(s) exactly as directed by your doctor. It is easier to prevent nausea than to treat it. • Contact your doctor if nausea lasts more than 48 hours or vomiting for more than 24 hours. • Also see Nausea & Vomiting pamphlet.* 	<p>Contact your health care team if no improvement or if severe</p>
<p><i>Mouth sores</i></p> <ul style="list-style-type: none"> • Maintain good mouth hygiene. Regular teeth brushing with a soft toothbrush or Toothette®, and regular use of alcohol-free mouthwashes. • Instead, try a homemade mouthwash: Mix 1 teaspoonful of baking soda and 1 teaspoonful of salt in 4 cups (1L) of water. • Avoid hot, spicy, acidic, hard or crunchy foods. • Check with your doctor or nurse as soon as you notice sores in mouth/lips or pain with swallowing. Your doctor may prescribe a prescription mouthwash to relieve mouth sores and prevent infection. • Also see Mouth Care pamphlet.* 	<p>Contact your health care team as soon as possible</p>
<p><i>Hair thinning or loss</i></p> <ul style="list-style-type: none"> • Use a gentle soft brush; care should be taken with hair sprays, bleaches, dyes and perms. • Your hair usually grows back after your treatment ends, but the texture or colour may change. 	
<p><i>Tiredness</i></p> <ul style="list-style-type: none"> • Rest often; take naps if needed. Move slowly when getting up. • Eat well-balanced meals and drink plenty of fluids. Light exercise may help. • Do not drive a motor vehicle or operate machinery when feeling tired. 	<p>Contact your health care team if no improvement or if severe</p>
<p><i>Poor Appetite; don't feel like eating; weight loss</i></p> <ul style="list-style-type: none"> • Eat foods that you like and try to eat regular small meals. • Use meal supplements if possible. See a dietitian. 	<p>Contact your health care team if no improvement or if severe</p>

*The most updated version and more symptom control information can be found on: <http://www.cancercare.on.ca/druginfo>

Prepared with input from the Cancer Care Ontario-Medication Information Sheets Working Group.

May 2018

pegylated liposomal DOXOrubicin

Side effects and what to do	When to contact doctor?
More Common Side Effects	
<p>Rash (rarely severe); dry, itchy skin</p> <ul style="list-style-type: none"> • Stay out of the sun; wear sunblock, a hat and cover exposed skin. • Use daily moisturizer. • If severe, get emergency medical help right away. 	<p>Contact your health care team if no improvement or if severe</p>
<p>Sensitivity reaction (usually the first injection only) (flushing, rash, itchiness, facial swelling, headache, chest or throat tightness)</p> <ul style="list-style-type: none"> • Watch for symptoms during and shortly after the infusion <p>Allergic reaction is less common (fever, severe rash, itchiness, swollen face, lip or tongue, chest or throat tightness)</p>	<p>Get emergency medical help right away</p>
<p>Unusual bleeding or bruising</p> <p>You may have black stools, cough up blood, blood in your urine, purple or red dots on your skin or bleeding that will not stop.</p> <p>Fever, chills, infection</p> <p>You have a fever if your temperature taken in your mouth (oral temperature) is:</p> <ul style="list-style-type: none"> • 38.3°C (100.9°F) or higher at any time OR • 38.0°C (or 100.4°F) or higher for at least one hour. <p>While you are getting chemotherapy treatments:</p> <ul style="list-style-type: none"> • Keep a digital thermometer at home and take your temperature if you feel hot or unwell (for example, chills). • Avoid taking medications that treat a fever before you take your temperature (for example, Tylenol®, acetaminophen, Advil® or ibuprofen) as they may hide a fever. • Do not eat or drink anything hot or cold right before taking your temperature. • Wash your hands often. • Check with your doctor before getting any vaccines, surgeries, medical procedures or visiting your dentist. <p>If you have a fever, talk to your health care team or go to the closest emergency room. See our Neutropenia (Low white blood cell count) pamphlet for more information.</p>	<p>Get emergency medical help right away</p>

*The most updated version and more symptom control information can be found on: <http://www.cancercare.on.ca/druginfo>
Prepared with input from the Cancer Care Ontario-Medication Information Sheets Working Group.
May 2018

pegylated liposomal DOXOrubicin

Side effects and what to do	When to contact doctor?
Less Common Side Effects, but may be Severe	
Blockage of an artery (blood vessel) in your heart, brain, chest, belly, or limbs; this may result in stroke (sudden loss of vision, speech, or the use of your limb(s)) or heart attack (chest pain, shortness of breath), or pain in chest, belly or limb	Get emergency medical help right away
Heart problems (irregular heartbeat, chest pain, fainting, swelling, shortness of breath)	Get emergency medical help right away
Pain, burning, redness, or swelling on skin where drug was injected <ul style="list-style-type: none"> • Let your healthcare team know right away when this happens, since this drug can harm or irritate tissues if it leaks from the vein during injection. 	
Blood clot (limb pain or swelling, hardened vein in limb), may occur in lungs (sudden start of coughing, breathing problems, chest pain, coughing blood)	Get emergency medical help right away

For more links on how to manage your symptoms go to www.cancercareontario.ca/symptoms.

The information set out in the medication information sheets, regimen information sheets, and symptom management information (for patients) contained in the Drug Formulary (the "Formulary") is intended to be used by health professionals and patients for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or side effects of a certain drug, nor should it be used to indicate that use of a particular drug is safe, appropriate or effective for a given condition.

A patient should always consult a healthcare provider if he/she has any questions regarding the information set out in the Formulary. The information in the Formulary is not intended to act as or replace medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.