

How to Manage Constipation: For People With Cancer

The recommendations in this guide are for people before, during and after cancer treatment.

If you spend most of your time in bed, are unable to eat or have cancer in your digestive tract, please speak to your health care team about your constipation before using this guide.

What is constipation?

Constipation is when you have bowel movements (go poo) less often than what is normal for you. Getting constipated is common for people with cancer.

You may be constipated if:

- Your stool (poo) is hard to pass
- You feel you need to have a bowel movement but nothing comes out
- Your stool is small, hard, dry, and looks like pellets

When you are constipated you may also have:

- A stomach ache or cramps
- A bloated belly that feels full or uncomfortable
- A lot of gas or burping
- Nausea (feeling like throwing up) or vomiting (actually throwing up)

How often should you have a bowel movement?

Not everyone has a bowel movement every day. If you are eating less, you may go less often. As long as your poo is soft and easy to pass, you are not constipated.



When should I talk to my health care team?

Contact your health care team right away if you have any of the symptoms below:

- You have not passed any stool (poo) in more than 3 days
- Bright red blood in your stool
- Very dark or black stool
- You are not able to pass gas
- Bloating, cramping or pain/discomfort in your stomach
- Nausea and vomiting (throwing up)

Risk of bowel obstruction

You may be at risk of a bowel obstruction (blocked intestines) if you have cancer in your abdomen (belly) or you have had bowel surgery. Bowel obstruction can feel similar to constipation. If you have a bowel obstruction you will not be able to pass any gas or poo. If you have had bowel surgery, tell your health care team right away or go to the nearest emergency room (ER) if you have any signs of constipation.

What causes constipation when you have cancer?

Medications are the most common cause of constipation when you have cancer. Some of the medications that cause constipation are:

- Opioid pain medications (like morphine)
- Chemotherapy medications (like vincristine, vinorelbine)
- Medicines for nausea (like ondansetron)
- Supplements (like calcium or iron medication)

Stool softeners (like soflox and docusate sodium) do not help most cases of constipation. Speak with your doctor or pharmacist if you are taking them.

Some other examples of things that may cause constipation are:

- Changes in your diet
- Not drinking enough liquids
- Stress, anxiety and depression
- Not getting enough exercise or physical activity

What can I do to manage constipation?

Take laxatives

Laxatives are medicines that help you have a bowel movement (go poo).

You may need to take laxatives:

- Every now and then for occasional constipation
- Regularly as part of a bowel care routine if you take medication that causes constipation

There are different kinds of laxatives that work in different ways. Your health care team can help you choose the type of laxative that is right for you.

Laxatives can cause mild stomach cramps. If you get painful cramps or diarrhea, speak to your doctor or pharmacist about whether you should stop taking laxatives.

What is a bowel care routine?

- A bowel care routine is a plan to manage constipation with laxatives
- Your cancer centre may have a bowel care routine for you to follow if you are taking opioid pain medications
- If you stop taking laxatives while taking opioids, you will get constipated

Other things that can help constipation

Drink enough liquids: Liquids help to make your stool soft and easy to pass. Drink at least 6 - 8 cups of liquids each day unless your health care team has told you to drink more or less. If drinking liquids is hard, take small sips often or eat watery fruit, such as melon, pineapple and plums.

Move your body. Any kind of exercise or movement may help to relieve constipation. Start slowly with light exercise and movement, such as walking. Even 10 minutes a day can help. Stop and rest if you feel pain, nausea or if you have trouble breathing.

Try to relax about using the bathroom. Feeling stress or anxiety about having to poo can make it harder to go. Take your time and try to relax.

Ask for privacy if you need to. If you can, use the bathroom or commode, instead of a bedpan.

Make a bathroom routine. Try to go to the bathroom around the same time each day. You may find it easier to have a bowel movement first thing in the morning or after a meal.

Get in a better position on the toilet. Put a low footstool under your feet when sitting on the toilet. Do not push hard or strain when trying to pass stool (poo).

Key Points

- 1) You may need to take laxatives to help manage your constipation
- 2) If you are taking an opioid medication, you may need a bowel care routine
- 3) Reach out to your health care team if you have any concerns or questions about managing your constipation

Where to get more information about constipation

The Canadian Cancer Society is a trusted source of information. Visit them at cancer.ca or phone 1-888-939-3333

For guides to help manage other symptoms please visit cancercareontario.ca/en/symptom-management.

The recommendations in this guide are based on published evidence and expert consensus.

This patient guide was created by Ontario Health (Cancer Care Ontario) to provide patients with information about constipation. It should be used for information only and does not replace medical advice. This guide does not include all information available about managing constipation. Always ask your health care team if you have questions or concerns. The information in this guide does not create a physician-patient relationship between Ontario Health (Cancer Care Ontario) and you.

Ontario Health (Cancer Care Ontario) retains all copyright, trademark, and all other rights in the guide, including all text and graphic images. No portion of this resource may be used or reproduced, other than for personal use, or distributed, transmitted or "mirrored" in any form, or by any means, without the prior written permission of Ontario Health (Cancer Care Ontario).

Need this information in an accessible format? 1-877-280-8538, TTY 1-800-855-0511, info@ontariohealth.ca

Updated March 2023